arXiv:1510.00145v1 [math.AP] 1 Oct 2015

Integral representation for functionals defined on SBD^p in dimension two

October 2, 2015

Sergio Conti[1], Matteo Focardi[2], and Flaviana Iurlano[1]

[1] *Institut für Angewandte Mathematik, Universität Bonn*
53115 Bonn, Germany

[2] *DiMaI, Università di Firenze*
50134 Firenze, Italy

We prove an integral representation result for functionals with growth conditions which give coercivity on the space $SBD^p(\Omega)$, for $\Omega \subset \mathbb{R}^2$. The space SBD^p of functions whose distributional strain is the sum of an L^p part and a bounded measure supported on a set of finite $\mathcal{H}^1$-dimensional measure appears naturally in the study of fracture and damage models. Our result is based on the construction of a local approximation by $W^{1,p}$ functions. We also obtain a generalization of Korn's inequality in the SBD^p setting.

1 Introduction

The direct methods of Γ-convergence are of paramount importance in studying variational limits and relaxation problems since their introduction in the seminal paper by Dal Maso and Modica [27]. They focus on the study of abstract limiting functionals $F(u, A)$, obtained for instance using $\overline{\Gamma}$-convergence arguments; one key ingredient is the proof of an integral representation for $F(u, A)$. Here $u : \Omega \to \mathbb{R}^N$ is an element of a suitable function space $\mathscr{X}(\Omega)$, and A runs in the class $\mathcal{A}(\Omega)$ of open subsets of a given open set $\Omega \subset \mathbb{R}^n$. The notion of *variational functional* is at the heart of the matter: F, regarded as depending on the couple $(u, A) \in \mathscr{X}(\Omega) \times \mathcal{A}(\Omega)$, has to satisfy suitable lower semicontinuity, locality and measure theoretic properties (for more details see properties (i)-(iii) in Theorem 1.1). The specific growth conditions of the functional determine the natural functional space in which the function u lies. Under these assumptions $F(u, A)$ can be written as an integral over the domain of integration A with respect to a suitable measure. The integrands may depend on x, $u(x)$ and $\nabla u(x)$, and possibly on other local quantities of u, such as higher order or distributional derivatives. Furthermore, as first shown in some cases in [28] and then generalized in [10], the corresponding

energy densities can be characterized in terms of cell formulas, i.e. asymptotic Dirichlet problems on small cubes or balls involving F itself, with boundary data depending on the local properties of u.

Integral representation results have been obtained in several contexts with increasing generality: starting with the pioneering contribution by De Giorgi for limits of area-type integrals [29], it has been extended to functionals defined first on Sobolev spaces in [38, 15, 12, 14, 13] and on the space of functions with Bounded Variation in [24, 8], and then to energies defined on partitions in [2] and on the subspace SBV in [11] (we refer to [13, 25, 10, 9] for a more exhaustive list of references). The global method for relaxation introduced and developed in [10, 9] provides a general approach that unifies and extends the quoted results.

We address the integral representation of functionals defined on the subspace $SBD^p(\Omega)$ of the space $BD(\Omega)$ in two dimensions. The space of functions of bounded deformation $BD(\Omega)$ is characterized by the fact that the symmetric part of the distributional gradient $Eu := (Du + Du^T)/2$ of $u \in L^1(\Omega, \mathbb{R}^n)$ is a bounded Radon measure, namely

$$BD(\Omega) := \{u \in L^1(\Omega; \mathbb{R}^n) : Eu \in \mathcal{M}(\Omega; \mathbb{R}^{n\times n}_{\rm sym})\},$$

where $\Omega \subseteq \mathbb{R}^n$ is an open set, see [39, 40, 4]. BD and its subspaces SBD and SBD^p constitute the natural setting for the study of plasticity, damage and fracture models in a geometrically linear framework [39, 40, 41, 6, 35]. In particular, SBD^p is the set of BD functions such that the strain Eu can be written as the sum of a function in $L^p(\Omega, \mathbb{R}^{n\times n})$ and a part concentrated on a rectifiable set with finite $\mathcal{H}^{n-1}$-measure, see [7, 16, 17, 19].

For functionals with linear growth defined on SBD an integral representation result was obtained by Ebobisse and Toader [31]. These functionals, however, lack coercivity on the relevant space. The situation of functionals defined on SBD^p and with corresponding growth properties is open. We give here a solution in two dimensions.

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^2$ be a bounded Lipschitz set, $p \in (1, \infty)$, $F : SBD^p(\Omega) \times \mathcal{B}(\Omega) \to [0, \infty)$ be such that*

(i) $F(u, \cdot)$ is a Borel measure for any $u \in SBD^p(\Omega)$;

(ii) $F(\cdot, A)$ is lower semicontinuous with respect to the strong $L^1(\Omega, \mathbb{R}^2)$-convergence for any open set $A \subset \Omega$;

(iii) $F(\cdot, A)$ is local for any open set $A \subset \Omega$;

(iv) There are $\alpha, \beta > 0$ such that for any $u \in SBD^p(\Omega)$, any $B \in \mathcal{B}(\Omega)$,

$$\begin{aligned}\alpha\Big(\int_B |e(u)|^p dx + \int_{J_u \cap B} (1 + |[u]|) d\mathcal{H}^1\Big) \le F(u, B)\\ \le \beta\Big(\int_B (|e(u)|^p + 1) dx + \int_{J_u \cap B} (1 + |[u]|) d\mathcal{H}^1\Big).\end{aligned} \tag{1.1}$$

Then there are two Borel functions $f : \Omega \times \mathbb{R}^2 \times \mathbb{R}^{2\times 2} \to [0, \infty)$ and $g : \Omega \times \mathbb{R}^2 \times \mathbb{R}^2 \times S^1 \to [0, \infty)$ such that

$$F(u, B) = \int_B f(x, u(x), \nabla u(x)) dx + \int_{B \cap J_u} g(x, u^-(x), u^+(x), \nu_u(x)) d\mathcal{H}^1 . \tag{1.2}$$

Above and throughout the paper we will refer to the book [5] and to the papers [4, 7] for the notation and results about BV and BD spaces, respectively. In particular, $\mathcal{B}(\Omega)$ is the family of Borel subsets of Ω.

The proof of Theorem 1.1, which is given in Section 4, follows the general strategy introduced in [10, 9]. Their approach was based on a Poincaré-type inequality in SBV by De Giorgi, Carriero and Leaci, which is not known in SBD^p (see [30, 5]). Our main new ingredient is the construction of an approximation by $W^{1,p}$ functions, discussed in Section 3, which permits to bypass the De Giorgi-Carriero-Leaci inequality. The approximation is done so that the function is only modified outside a countable set of balls with small area and perimeter. In each ball, we give a construction of a $W^{1,p}$ extension for the SBD^p function by constructing a finite-element approximation on a countable mesh, which is chosen depending on the function u, see Section 2.

Our $W^{1,p}$ approximation result also leads naturally to the proof of the following variant of Korn's inequality for SBD^p functions.

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^2$ be a connected, bounded, Lipschitz set and let $p \in (1, \infty)$. Then there exists a constant c, depending on p and Ω, with the following property: for every $u \in SBD^p(\Omega)$ there exist a set $\omega \subset \Omega$ of finite perimeter, with $\mathcal{H}^1(\partial\omega) \le c\mathcal{H}^1(J_u)$, and an affine function $a(x) = Ax + b$, with $A \in \mathbb{R}^{2\times 2}$ skew-symmetric and $b \in \mathbb{R}^2$, such that*

$$\begin{aligned}\|u - a\|_{L^p(\Omega\setminus\omega, \mathbb{R}^2)} &\le c\|e(u)\|_{L^p(\Omega, \mathbb{R}^{2\times 2})},\\ \|\nabla u - A\|_{L^p(\Omega\setminus\omega, \mathbb{R}^{2\times 2})} &\le c\|e(u)\|_{L^p(\Omega, \mathbb{R}^{2\times 2})}.\end{aligned}$$

This improves a result of [33] to the sharp exponent. Variants of the first inequality were first obtained in [18, 32].

The construction of Section 3 turns out to be crucial also in proving existence for the Griffith fracture model, generalizing [30] to the case of linearized elasticity in dimension two. This will be the object of the forthcoming paper [20].

2 Approximation of SBD^p functions with small jump set

In this Section we prove the following approximation result.

Theorem 2.1. *Let $n = 2$, $p \in [1, \infty)$. There exists $\eta > 0$ and $\tilde{c} > 0$ such that if $J \in \mathcal{B}(B_{2r})$, for some $r > 0$, satisfies*

$$\mathcal{H}^1(J) < 2r\eta, \tag{2.1}$$

then there exist $R \in (r, 2r)$ for which the following holds: for every $u \in SBD^p(B_{2r})$ with $\mathcal{H}^1(J_u \cap B_{2r} \setminus J) = 0$ there exists $\phi(u) \in SBD^p(B_{2r}) \cap W^{1,p}(B_R, \mathbb{R}^2)$ such that

(i) $\mathcal{H}^1(J_u \cap \partial B_R) = 0$;

(ii) $\displaystyle\int_{B_R} |e(\phi(u))|^q dx \le \tilde{c} \int_{B_R} |e(u)|^q dx$, *for every* $q \in [1, p]$;

(iii) $\|u - \phi(u)\|_{L^1(B_R, \mathbb{R}^2)} \le \tilde{c} R |Eu|(B_R)$;

(iv) $u = \phi(u)$ *on* $B_{2r} \setminus B_R$, $\mathcal{H}^1(J_{\phi(u)} \cap \partial B_R) = 0$;

(v) if $u \in L^\infty(B_{2r}, \mathbb{R}^2)$, *then* $\|\phi(u)\|_{L^\infty(B_{2r}, \mathbb{R}^2)} \le \sqrt{2} \|u\|_{L^\infty(B_{2r}, \mathbb{R}^2)}$.

Proof. We follow an idea of [23, 22].

Arguing as in [23, Lemma 4.3] we first claim that there exists $R \in (r, 2r)$ such that for $\delta_k := R\, 2^{-k}$ we have

$$\mathcal{H}^1(J \cap \partial B_R) = 0, \tag{2.2}$$

$$\mathcal{H}^1(J \cap (B_R \setminus B_{R-\delta_k})) < 20\eta\delta_k, \quad \text{for every } k \in \mathbb{N}. \tag{2.3}$$

To prove this, we first observe that (2.2) holds for almost every R, therefore it suffices to show that (2.3) holds on a set of positive measure. We consider the family of intervals

$$\{[R - \delta_k, R] : \mathcal{H}^1(J \cap (B_R \setminus B_{R-\delta_k})) \ge 20\eta\delta_k\}$$

and we define I as the union of all intervals of the family, with $R \in (r, 2r)$, $k \in \mathbb{N}$. By Vitali's covering theorem, there exists a countable set $(R_i, k_i)_{i \in \mathbb{N}}$ such that the corresponding intervals $[R_i - \delta_{k_i}, R_i]$ are pairwise disjoint and cover at least one fifth of I. Therefore by (2.1) we obtain

$$2r\eta > \mathcal{H}^1(J \cap B_{2r}) \ge \sum_{i \in \mathbb{N}} \mathcal{H}^1(J \cap (B_{R_i} \setminus B_{R_i - \delta_{k_i}})) \ge \sum_{i \in \mathbb{N}} 20\eta\delta_{k_i}.$$




Figure 1: Sketch of the construction of the grid in the proof of Theorem 2.1.

Since $\delta_{k_i} = \mathcal{L}^1([R_i - \delta_{k_i}, R_i])$, we conclude that $\mathcal{L}^1(I) < r$. This proves the existence of R such that (2.2) and (2.3) hold, which is fixed for the rest of the proof.

We define $R_k := R - \delta_k$ and $\overline{x}_{k,j} := R_k(\cos\frac{2\pi j}{2^k}, \sin\frac{2\pi j}{2^k})$, $j = 1, \dots, 2^k$. We say that $\overline{x}_{k,j}$ and $\overline{x}_{k',j'}$ are neighbors if either $k = k'$ and $j = j' \pm 1$, working modulo 2^k, or (up to a permutation) $k = k' + 1$ and $j \in \{2j' - 1, 2j', 2j' + 1\}$, again modulo 2^k. This gives a decomposition of B_R into countably many triangles, whose angles are uniformly bounded away from 0 and π, see Figure 1.

We shall construct $\phi(u)$ as a linear interpolation on a triangulation whose vertices are slight modifications of $\overline{x}_{k,j}$. Following the idea of [22, Proposition 2.2], we next show how to construct the modified triangulation. We start off considering two neighboring points $\overline{x}$ and $\overline{y}$ in $\{\overline{x}_{k,j}\}_{k,j}$, connected by the segment $S_{\overline{x},\overline{y}} \subset \overline{B}_{R_{k+1}} \setminus B_{R_{k-1}}$ for some k, and notice that $c_1\delta_k \le |\overline{x} - \overline{y}| \le c_2\delta_k$ for some $c_1 \in (0,1)$, $c_2 > 1$ independent from k. Let $\alpha := c_1/(8c_2)$ and consider the convex envelope

$$O_{\overline{x},\overline{y}} := \mathrm{conv}\,(B(\overline{x}, \alpha\delta_k) \cup B(\overline{y}, \alpha\delta_k)). \tag{2.4}$$

Let $a_{\overline{x},\overline{y}}$ denote the infinitesimal rigid movement appearing in the Poincaré's inequality for u on the set

$$Q_{\overline{x},\overline{y}} := \{\xi \in B_R : \mathrm{dist}(\xi, S_{\overline{x},\overline{y}}) < |\overline{x} - \overline{y}|/(8c_2)\}.$$

Given $\vartheta \in (0,1)$, let us prove that for η sufficiently small and $\tilde{c}$ sufficiently large, depending only on ϑ, there exists a subset $F \subset B(\overline{x}, \alpha\delta_k) \times B(\overline{y}, \alpha\delta_k)$ with $\frac{(\mathcal{L}^2 \times \mathcal{L}^2)(F)}{\mathcal{L}^2(B_{\alpha\delta_k})^2} < \vartheta$, such that for every $(x,y) \notin F$ the one-dimensional section u_z^ν has the following properties:




Figure 2: Slice along the line of direction $\nu = (x-y)/|x-y|$ passing through z in the proof of Theorem 2.1.

(P1) $u_z^\nu \in SBV(s_{x,y})$;

(P2) $\mathcal{H}^0(J_{u_z^\nu}) = 0$, so that $u_z^\nu \in W^{1,1}(s_{x,y})$;

(P3) $\displaystyle\int_{s_{x,y}} |(u_z^\nu)'|dt \le \frac{\tilde{c}}{\delta_k} \int_{O_{\overline{x},\overline{y}}} |e(u)|dx'$;

(P4) $|u(\xi) - a_{\overline{x},\overline{y}}(\xi)| \le \dfrac{\tilde{c}}{\delta_k} |Eu|(Q_{\overline{x},\overline{y}})$, for $\xi = x, y$.

Here $u_z^\nu(t) := u(z + t\nu) \cdot \nu$ is the slice of u along the line of direction

$$\nu := \frac{x-y}{|x-y|}, \tag{2.5}$$

and passing through

$$z := (\mathrm{Id} - \nu \otimes \nu)x \in \mathbb{R}\nu^\perp \cap (x + \mathbb{R}\nu) \tag{2.6}$$

where $\mathbb{R}\nu^\perp$ is the linear space orthogonal to ν, and $s_{x,y} \subset \mathbb{R}$ is the segment such that $z + s_{x,y}\nu = S_{x,y}$, see Figure 2.

In order to obtain property (P1) we first define the measure $\mu_{\nu,z} := |u_z^\nu|\mathcal{L}^1$ and we observe that

$$\begin{aligned} &\int_{B(\overline{x},\alpha\delta_k)\times B(\overline{y},\alpha\delta_k)} \mu_{\nu,z}(s_{x,y})dx\,dy \\ &\qquad \le c\delta_k^2 \int_{B(\overline{x},\alpha\delta_k)} dx \int_{\{|\nu-\overline{\nu}|\le c(\alpha)\}} \mu_{\nu,z}((O_{\overline{x},\overline{y}})_z^\nu) d\mathcal{H}^1(\nu), \end{aligned} \tag{2.7}$$

by the change of variables $y = x + s\nu$, where $\overline{\nu}$ is defined as ν with $\overline{x}, \overline{y}$ in place of x, y and $(O_{\overline{x},\overline{y}})^\nu_z \subset \mathbb{R}$ corresponds to $O_{\overline{x},\overline{y}} \cap (z + \mathbb{R}\nu)$. By Fubini's theorem the last term in the previous inequality is less than or equal to

$$c\delta_k^3 \int_{\{|\nu-\overline{\nu}|\le c(\alpha)\}} d\mathcal{H}^1(\nu) \int_{\nu^\perp} \mu_{\nu,z}((O_{\overline{x},\overline{y}})^\nu_z) d\mathcal{H}^1(z) \le c\delta_k^3 \mu(O_{\overline{x},\overline{y}}), \tag{2.8}$$

where $\mu(O_{\overline{x},\overline{y}}) := \int_{O_{\overline{x},\overline{y}}} |u| dx'$. By (2.7) and (2.8) the set $F_1 \subset B(\overline{x}, \alpha\delta_k) \times B(\overline{y}, \alpha\delta_k)$ of points (x, y), for which the inequality

$$\mu_{\nu,z}(s_{x,y}) > \frac{\tilde{c}}{\delta_k} \mu(O_{\overline{x},\overline{y}})$$

holds, satisfies $\frac{(\mathcal{L}^2 \times \mathcal{L}^2)(F_1)}{\mathcal{L}^2(B_{\alpha\delta_k})^2} < \vartheta/16$, for $\tilde{c}$ large enough, depending on ϑ and α.

For $(x, y) \in B(\overline{x}, \alpha\delta_k) \times B(\overline{y}, \alpha\delta_k) \setminus F_1$ we now repeat the argument in (2.7) and (2.8) above redefining

$$\mu_{\nu,z} := |D(u^\nu_z)|.$$

We find that for (x, y) out of a small (in the previous sense) set F_1' one has $u^\nu_z \in SBV(s_{x,y})$ and

$$|D(u^\nu_z)|(s_{x,y}) \le \frac{\tilde{c}}{\delta_k} |Eu|(O_{\overline{x},\overline{y}}).$$

As for property (P2), we use (2.7) and (2.8) with $\mu_{\nu,z} := \mathcal{H}^0 \llcorner (J_{u^\nu_z} \cap s_{x,y})$ and $\mu := \mathcal{H}^1 \llcorner (J_u \cap O_{\overline{x},\overline{y}})$. Now (2.1) implies

$$\int_{B(\overline{x},\alpha\delta_k) \times B(\overline{y},\alpha\delta_k)} \mathcal{H}^0(J_{u^\nu_z} \cap s_{x,y}) dx\, dy \le c\delta_k^4 \eta,$$

and hence the set F_2 of points (x, y) for which $\mathcal{H}^0(J_{u^\nu_z} \cap s_{x,y}) > 1/2$ is also small in the previous sense, for η small enough. Note that this is the only step which requires the hypothesis on the dimension $n = 2$.

Analogously properties (P3) and (P4) can be derived. From the argument above it is straightforward that for many points $x \in B(\overline{x}, \alpha\delta_k)$, still in the sense of a large ϑ-fraction of $B(\overline{x}, \alpha\delta_k)$, there are many points $y \in B(\overline{y}, \alpha\delta_k)$ for which $(x, y) \notin F$.

Let us construct now the modified grid with an iterative process (see also [22, Proposition 3.4]). We will use the notation B_i to indicate the balls $B(\overline{x}_{k,j}, \alpha\delta_k)$, lexicographically ordered.

We start by fixing a point $x_0 \in B_0$ for which there are many good choices in each neighboring ball. We next select $x_1 \in B_1$ among the points which are

good choices for x_0 and which have many good choices in each neighboring subsequent ball B_i, $i \geq 2$. Iterating the process, the point $x_m \in B_m$ will be taken among the good choices for the neighboring previously fixed x_i, $i < m$, and with the property that have many good choices in the neighboring subsequent B_i, $i > m$. Since each ball can have at most seven neighbors, at each step we select x_m avoiding just a small subset of B_m.

We call S the set of points obtained by this process and we construct a new triangulation, with x, y neighbors if and only if $\bar{x}$, $\bar{y}$ are neighbors. Notice that again

$$c_1\delta_k \leq |x - y| \leq c_2\delta_k, \tag{2.9}$$

for every couple of neighboring points x, y, with the same k as for the corresponding reference points $\overline{x}$ and $\overline{y}$, and suitable $c_1, c_2 > 0$ independent from k. We finally define $\phi(u)$ as the linear interpolation between the values of $u(x)$, $x \in S$ on each triangle of the triangulation.

Fixed a triangle T and any couple of its vertices x, y, we compute a component of the constant matrix $e(\phi(u))$ on T by

$$e(\phi(u))\nu \cdot \nu = \frac{(\phi(u)(x) - \phi(u)(y)) \cdot \nu}{|x - y|} = \fint_{s_{x,y}} (u_z^\nu)' dt, \tag{2.10}$$

where ν and z are defined in (2.5) and (2.6). We used the fact that u and $\phi(u)$ agree on x and y and that u is $W^{1,1}(s_{x,y})$ by the choice of x and y. By (2.10), (2.9), and property (P3) above it follows

$$|e(\phi(u))\nu \cdot \nu| \leq \frac{\tilde{c}}{\delta_k^2} \int_C |e(u)| dx',$$

where C is defined in (2.4). We recall that here and henceforth $\tilde{c}$ can possibly change. Letting ν vary among the directions of the sides of T, we obtain a control on the whole $|e(\phi(u))|$ thanks to (2.9)

$$|e(\phi(u))| \leq \frac{\tilde{c}}{\delta_k^2} \int_{C_T} |e(u)| dx', \tag{2.11}$$

where C_T denotes the convex envelope

$$C_T := \text{conv}\left(\cup B(\overline{x}, \alpha\delta_k)\right)$$

and the union is taken over the three vertices $\overline{x}$ in the old triangulation corresponding to the three vertices of T. We remark that $B(\overline{x}, \alpha\delta_k) \subset B_{R_{k+1}} \setminus B_{R_{k-1}}$ for all $\overline{x} \in \partial B_{R_k}$, therefore the sets C_T have finite overlap.

We are ready to prove property (ii). By Jensen's inequality and (2.11) we have for $1 \le q \le p$

$$\int_T |e(\phi(u))|^q dx' = \mathcal{L}^2(T)|e(\phi(u))|^q \le \tilde{c}\mathcal{L}^2(C_T)\Big(\fint_{C_T} |e(u)|dx'\Big)^q$$
$$\le \tilde{c}\int_{C_T} |e(u)|^q dx',$$

and finally summing up on all triangles T we get the conclusion.

In order to prove properties (iii) and (iv) we estimate

$$\int_T |u-\phi(u)|dx' \le \int_T |u - a_{\overline{x},\overline{y}}|dx' + \int_T |a_{\overline{x},\overline{y}} - \phi(u)|dx', \tag{2.12}$$

where T is again a triangle of the modified triangulation with vertices x, y, z, $\overline{x}, \overline{y}, \overline{z}$ denote the three corresponding vertices of the old triangulation, $a_{\overline{x},\overline{y}}$ is the infinitesimal rigid motion appearing in the Poincaré's inequality for u on $Q_{\overline{x},\overline{y}}$ (see item (P4) above).

Let us study first the second term in (2.12). Since $a_{\overline{x},\overline{y}} - \phi(u)$ is affine, it achieves its maximum on a vertex ξ of T, therefore

$$\int_T |a_{\overline{x},\overline{y}} - \phi(u)|dx' \le c\delta_k^2 |a_{\overline{x},\overline{y}}(\xi) - \phi(u)(\xi)| = c\delta_k^2 |a_{\overline{x},\overline{y}}(\xi) - u(\xi)|.$$

Notice that if $\xi = \overline{z}$ then

$$\begin{aligned} c\delta_k^2|a_{\overline{x},\overline{y}}(\xi) - u(\xi)| &\le c\delta_k^2|a_{\overline{x},\overline{y}}(\xi) - a_{\overline{x},\xi}(\xi)| + c\delta_k^2|a_{\overline{x},\xi}(\xi) - u(\xi)| \\ &\le c\int_{B(\overline{x},\alpha\delta_k)} |a_{\overline{x},\overline{y}} - a_{\overline{x},\xi}|dx' + c\delta_k|Eu|(Q_{\overline{x},\xi}), \end{aligned} \tag{2.13}$$

where we used the fact that $a_{\overline{x},\xi}, a_{\overline{x},\overline{y}}$ are affine and item (P4) above; if $\xi \in \{\overline{x}, \overline{y}\}$ then only the second term appears.

By (2.12)-(2.13), the triangular inequality, and Poincaré's inequality we conclude

$$\int_T |u - \phi(u)|dx' \le c\delta_k |Eu|(Q_T), \tag{2.14}$$

where $Q_T := Q_{\overline{x},\overline{y}} \cup Q_{\overline{y},\overline{z}} \cup Q_{\overline{z},\overline{x}}$. Finally summing up over T we obtain property (iii).

We prove now property (iv), property (v) holding true by construction. We define $\phi(u) := u$ outside $\overline{B}_R$ and know that $\phi(u) \in W^{1,p}(B_R, \mathbb{R}^2) \cap SBD(B_{2r})$. It remains to prove that the traces on ∂B_R coincide, or, equivalently, that $\mathcal{H}^1(J_{\phi(u)} \cap B_R) = 0$. Let $\psi_k \in C^\infty(B_R)$ be such that $\psi_k = 0$ on B_{R_k}, $\psi_k = 1$ in a neighborhood of ∂B_R, and $|\nabla\psi_k| \le c/\delta_k$. We define

$v_k := (u - \phi(u))\psi_k \in SBD(B_R)$ and we prove that $v_k \to 0$ strongly in BD, this implying in turn $v_k|_{\partial B_R} \to 0$ in $L^1(\partial B_R, \mathbb{R}^2)$ in the sense of traces and therefore property (iv). Clearly

$$\int_{B_R} |v_k| dx \leq \int_{B_R \setminus B_{R_k}} |u - \phi(u)| dx \to 0$$

by the dominated convergence theorem. Finally, using (2.14) and the fact that the triangles have finite overlap,

$$\begin{aligned} |Ev_k|(B_R) &\leq |E(u - \phi(u))|(B_R \setminus B_{R_k}) + \frac{c}{\delta_k} \int_{B_R \setminus B_{R_k}} |u - \phi(u)| dx \\ &\leq \tilde{c}|E(u - \phi(u))|(B_R \setminus B_{R_k}), \end{aligned}$$

the last term tends to 0 and this concludes the proof of property (iv).

□

3 Regularity of SBD^p functions with small jump set

We first discuss how SBD^p functions can be approximated by $W^{1,p}$ functions locally away from the jump set (Section 3.1), and then how they can be approximated by piecewise $W^{1,p}$ functions around the jump set (Section 3.3). Our approximation result also leads to the Korn inequality stated in Theorem 1.2. The key ingredient for all these results is the construction of Theorem 2.1. Throughout the section $\eta \in (0, 1)$ will be the constant from Theorem 2.1 and $n = 2$.

3.1 Approximation of SBD^p functions with $W^{1,p}$ functions

We shall use that the construction of Theorem 2.1, using a suitable covering argument, permits to approximate SBD^p functions by $W^{1,p}$ functions which coincide away from a small neighborhood of the jump set. The neighborhood is the union of countably many balls, such that each of them contains an amount of jump set proportional to the radius. Before discussing the covering argument in Proposition 3.2, we show that (away from the boundary) almost any point of the jump set is the center of a ball with the appropriate density.

Lemma 3.1. *Let $s \in (0,1)$. Let $J \in \mathcal{B}(B_{2\rho})$, for some $\rho > 0$, be such that $\mathcal{H}^1(J) < \eta(1-s)\rho$ then for $\mathcal{H}^1$-a.e. $x \in J \cap B_{2s\rho}$ there exists a radius $r_x \in (0,(1-s)\rho)$ such that*

$$\mathcal{H}^1\big(J \cap \partial B_{r_x}(x)\big) = 0, \tag{3.1}$$

$$\eta\, r_x \le \mathcal{H}^1\big(J \cap B_{r_x}(x)\big) \le \mathcal{H}^1\big(J \cap B_{2r_x}(x)\big) < 2\,\eta\, r_x. \tag{3.2}$$

Proof. We fix $x \in J \cap B_{2s\rho}$, choose $\lambda_x \in (\rho, 2\rho)$ such that $\mathcal{H}^1\big(J \cap \partial B_{\lambda_x/2^k}(x)\big) = 0$ for all $k \in \mathbb{N}$, and define

$$r_x := \max\{{}^{\lambda_x}\!/_{2^k} :\ k \in \mathbb{N},\ \mathcal{H}^1\big(J \cap B_{\lambda_x/2^k}(x)\big) \ge \eta\,\lambda_x 2^{-k}\}.$$

The set is nonempty for $\mathcal{H}^1$-almost every x because $\eta < 1$. The estimates (3.2) hold by definition. To conclude that $r_x < (1-s)\rho$ it is enough to notice that the opposite inequality would give the ensuing contradiction

$$\mathcal{H}^1(J) \ge \mathcal{H}^1\big(J \cap B_{r_x}(x)\big) \ge \eta\, r_x \ge (1-s)\eta\,\rho > \mathcal{H}^1(J).$$

□

We are now ready to prove the main result of the section via a covering argument, Lemma 3.1, and Theorem 2.1.

Proposition 3.2. *Let $p \in (1,\infty)$, $n = 2$. There exists a universal constant $c > 0$ such that if $u \in SBD^p(B_{2\rho})$, $\rho > 0$, satisfies*

$$\mathcal{H}^1(J_u \cap B_{2\rho}) < \eta\,(1-s)\rho$$

for $\eta \in (0,1)$ as in Theorem 2.1 and some $s \in (0,1)$, then there is a countable family $\mathcal{F} = \{B\}$ of closed balls of radius $r_B < (1-s)\rho$, each contained in $B_{(1+s)\rho}$, and a field $w \in SBD^p(B_{2\rho})$ such that

(i) $\rho^{-1}\sum_{\mathcal{F}} \mathcal{L}^2\big(B\big) + \sum_{\mathcal{F}} \mathcal{H}^1\big(\partial B\big) \le {}^{c}\!/_{\eta}\, \mathcal{H}^1(J_u \cap B_{2\rho})$;

(ii) $\mathcal{H}^1\big(J_u \cap \cup_{\mathcal{F}} \partial B\big) = \mathcal{H}^1\big((J_u \cap B_{2s\rho}) \setminus \cup_{\mathcal{F}} B\big) = 0$;

(iii) $w = u$ $\mathcal{L}^2$-a.e. on $B_{2\rho} \setminus \cup_{\mathcal{F}} B$;

(iv) $w \in W^{1,p}(B_{2s\rho}, \mathbb{R}^2)$ and $\mathcal{H}^1(J_w \setminus J_u) = 0$;

(v)

$$\int_{\cup_{\mathcal{F}} B} |e(w)|^p dx \le c \int_{\cup_{\mathcal{F}} B} |e(u)|^p dx; \tag{3.3}$$

and there exists a skew-symmetric matrix A such that

$$\int_{B_{2s\rho} \setminus \cup_{\mathcal{F}} B} |\nabla u - A|^p dx \le c(p) \int_{B_{2\rho}} |e(u)|^p dx; \tag{3.4}$$

(vi) $\|u-w\|_{L^1(B,\mathbb{R}^2)} \le c\, r_B\, |Eu|(B)$, *for every* $B \in \mathcal{F}$;

(vii) if, additionally, $u \in L^\infty(B_{2\rho},\mathbb{R}^2)$ *then* $w \in L^\infty(B_{2\rho},\mathbb{R}^2)$ *with*

$$\|w\|_{L^\infty(B_{2\rho},\mathbb{R}^2)} \le c\|u\|_{L^\infty(B_{2\rho},\mathbb{R}^2)}.$$

Proof. By Lemma 3.1 we find a family $\mathcal{F}'$ of open balls covering $\mathcal{H}^1$-a.e. $J_u \cap B_{2s\rho}$ that satisfies (3.1) and (3.2). Setting $J = J_u$, to every $B \in \mathcal{F}'$ we associate a new ball $B^* \subset B$ with the properties (i)-(v) of Theorem 2.1. Let $\mathcal{F}^*$ be the family of the new balls B^*, this is still a cover of J. Further, the balls B^* can be taken to be closed. By the Besicovitch covering theorem [5, Theorem 2.17] there are ξ countable subfamilies $\mathcal{F}'_j = \{B^i_j\}_{i\in\mathbb{N}}$ of disjoint balls. Therefore, setting $\mathcal{F} := \cup_{j=1}^{\xi}\mathcal{F}'_j$ we have $\mathcal{H}^1\big((J_u \cap B_{2s\rho}) \setminus \cup_{\mathcal{F}} B\big) = 0$. In addition, by (3.1) the first condition in item (ii) is satisfied as well, so that (ii) is established. Furthermore,

$$\begin{aligned}\sum_{B\in\mathcal{F}} \mathcal{H}^1(\partial B) =& 2\pi \sum_{B\in\mathcal{F}} r_B \overset{(3.2)}{\le} \frac{2\pi}{\eta}\sum_{B\in\mathcal{F}} \mathcal{H}^1\big(J_u \cap B\big)\\ \le& \xi\,\frac{2\pi}{\eta}\mathcal{H}^1\big(J_u \cap \cup_{B\in\mathcal{F}} B\big) \le \xi\,\frac{2\pi}{\eta}\mathcal{H}^1(J_u \cap B_{2\rho}).\end{aligned}$$

The volume estimate follows since $r_B \le \rho$ implies $\sum r_B^2 \le \rho \sum r_B$. We remark that a quadratic volume estimate also follows by $\sum r_B^2 \le (\sum r_B)^2$.

Let $\phi(u)$ be the function given by Theorem 2.1 on the balls of the first family $\mathcal{F}'_1$ and define for every $h \in \mathbb{N}$ a function

$$w_1^h := \begin{cases} \phi(u) & B_1^i,\ i \le h \\ u & \text{otherwise}\end{cases}$$

such that $w_1^h \in SBD^p(B_{2\rho})$, $w_1^h \in W^{1,p}(\cup_{i\le h} B_1^i;\mathbb{R}^2)$ with $w_1^h = u$ $\mathcal{L}^2$-a.e. on $B_{2\rho} \setminus \cup_{i\le h} B_1^i$ and $\mathcal{H}^1(J_{w_1^h} \setminus J_u) = 0$. In addition by item (ii) in Theorem 2.1

$$\begin{aligned}&\int_{B_{2\rho}} |e(w_1^h)|^p\,dx = \int_{\cup_{i\le h} B_1^i} |e(\phi(u))|^p\,dx + \int_{B_{2\rho}\setminus\cup_{i\le h} B_1^i} |e(u)|^p\,dx\\ &\le \tilde{c}\int_{\cup_{i\le h} B_1^i} |e(u)|^p\,dx + \int_{B_{2\rho}\setminus\cup_{i\le h} B_1^i} |e(u)|^p\,dx \le (1+\tilde{c})\int_{B_{2\rho}} |e(u)|^p\,dx, \quad (3.5)\end{aligned}$$

and

$$|Ew_1^h|(B_{2\rho}) \le |Eu|\big(B_{2\rho} \setminus \cup_{i\le h} B_1^i\big) + \tilde{c}\int_{\cup_{i\le h} B_1^i} |e(u)|\,dx.$$

Moreover, recalling that the B_1^i's are disjoint and that $w_1^{h-1} = u$ on B_1^h, item (iii) in Theorem 2.1 gives

$$\|w_1^h - w_1^{h-1}\|_{L^1(B_{2\rho};\mathbb{R}^2)} = \|w_1^h - u\|_{L^1(B_1^h;\mathbb{R}^2)} \le c\,\rho\,|Eu|(B_1^h),$$

in turn implying that for all $h \ge k \ge 1$

$$\|w_1^h - w_1^k\|_{L^1(B_{2\rho};\mathbb{R}^2)} \le \sum_{i=k+1}^{h} \|w_1^i - w_1^{i-1}\|_{L^1(B_1^h;\mathbb{R}^2)} \le c\,\rho\,|Eu|\big(\cup_{k+1\le i\le h} B_1^i\big).$$

Thus, $w_1^h \to w_1$ in $L^1(B_{2\rho};\mathbb{R}^2)$ with

$$w_1 := \begin{cases} \phi(u) & \cup_{\mathcal{F}_1'} B \\ u & \text{otherwise.} \end{cases}$$

The BD compactness theorem then yields that $w_1 \in BD(B_{2\rho})$. In turn, by (3.5) and since $\mathcal{H}^1(J_{w_1^h} \setminus J_u) = 0$, the SBD compactness theorem implies that actually $w_1 \in SBD^p(B_{2\rho})$ (see also [26, Theorem 11.3]). Furthermore, since

$$\mathcal{H}^1\big(J_{w_1^h} \cap \cup_{\mathcal{F}_1'} B\big) = \mathcal{H}^1(J_u \cap \cup_{i\ge h+1} B_1^i),$$

we may conclude that

$$\mathcal{H}^1\big(J_{w_1} \cap \cup_{\mathcal{F}_1'} B\big) \le \liminf_h \mathcal{H}^1\big(J_{w_1^h} \cap \cup_{\mathcal{F}_1'} B\big) = 0,$$

and therefore $w_1 \in W^{1,p}(\cup_{\mathcal{F}_1'} B, \mathbb{R}^2)$. Finally, by construction $w_1 = u$ $\mathcal{L}^2$-a.e. on $B_{2\rho} \setminus \cup_{\mathcal{F}_1'} B$ and $\mathcal{H}^1(J_{w_1} \setminus J_u) = 0$.

By iterating the latter construction, for all $1 < k \le \xi$ and for every $h \in \mathbb{N}$ we find

$$w_k^h := \begin{cases} \phi(w_{k-1}) & B_k^i,\ i \le h \\ w_{k-1} & \text{otherwise} \end{cases}$$

such that $w_k^h \in SBD^p(B_{2\rho})$, $w_k^h \in W^{1,p}(\cup_{i\le h} B_k^i;\mathbb{R}^2)$, $w_k^h = w_{k-1}$ $\mathcal{L}^2$-a.e. on $B_{2\rho} \setminus \cup_{i\le h} B_k^i$, $\mathcal{H}^1(J_{w_k^h} \setminus J_{w_{k-1}}) = 0$. In addition, arguing as above, $w_k^h \to w_k$ in $L^1(B_{2\rho},\mathbb{R}^2)$ with

$$w_k := \begin{cases} \phi(w_{k-1}) & \cup_{\mathcal{F}_k'} B \\ w_{k-1} & \text{otherwise,} \end{cases}$$

$w_k \in SBD^p(B_{2\rho})$, $w_k \in W^{1,p}(\cup_{j\le k} \cup_{\mathcal{F}_j'} B;\mathbb{R}^2)$, $w_k = w_{k-1}$ $\mathcal{L}^2$-a.e. on $B_{2\rho} \setminus \cup_{\mathcal{F}_k'} B$ and $\mathcal{H}^1(J_{w_k} \setminus J_{w_{k-1}}) = 0$.

Set $w := w_\xi$, then $w \in SBD^p(B_{2\rho})$, $w \in W^{1,p}(\cup_{\mathcal{F}} B; \mathbb{R}^2)$, $w = u$ $\mathcal{L}^2$-a.e. on $B_{2\rho} \setminus \cup_{\mathcal{F}} B$, $\mathcal{H}^1(J_w \setminus J_u) = 0$. Iterating estimate (3.5), inequality (3.3) follows at once with $c := \max\{1 + \tilde{c}, 2\pi\}\,\xi$, with $\tilde{c}$ the constant in Theorem 2.1. Korn's inequality (3.4) follows now immediately by (iii), (iv), and (3.3).

Finally, it is clear that also items (vi) and (vii) are satisfied in view of properties (iii) and (v) in Theorem 2.1. □

3.2 Korn's inequality in SBD^p

Proof of Theorem 1.2. By standard scaling and covering arguments it suffices to prove the assertion for a special Lipschitz domain. Precisely, let $\varphi : \mathbb{R} \to \mathbb{R}$ Lipschitz with $\min \varphi[(-1,1)] = 2$, and set $U := \{x : x_1 \in (-2,2), x_2 \in (-2, \varphi(x_1))\}$, and $U^{\rm int} := \{x : x_1 \in (-1,1), x_2 \in (-1, \varphi(x_1))\}$. It suffices to show that for any $u \in SBD^p(U)$ there are ω with $\mathcal{H}^1(\partial\omega) + |\omega|^{1/2} \leq c\mathcal{H}^1(J_u)$ and an affine function $a : \mathbb{R}^2 \to \mathbb{R}^2$ such that $\|u - a\|_{L^p(U^{\rm int}\setminus\omega, \mathbb{R}^2)} + \|\nabla u - \nabla a\|_{L^p(U^{\rm int}\setminus\omega, \mathbb{R}^2)} \leq c_{L,p} \|e(u)\|_{L^p(U, \mathbb{R}^{2\times 2})}$, with c depending on p and the Lipschitz constant L of φ. Obviously we can assume $\mathcal{H}^1(J_u)$ to be small.

Let $q_j := x_j + (-r_j/2, r_j/2)^2$ and $Q_j := x_j + (-r_j, r_j)^2$, and assume that $u \in SBD^p(Q_j)$ obeys $\mathcal{H}^1(J_u \cap Q_j) \leq \eta r_j/8$. By Proposition 3.2 with $\rho := r_j/2$ and $s := 1/\sqrt{2}$ and Poincaré's inequality there are ω_j and a_j affine with $r_j \mathcal{H}^1(\partial\omega_j) + |\omega_j|^{1/2} \leq c\mathcal{H}^1(J_u \cap Q_j)$ and $r_j^{-1}\|u_j - a_j\|_{L^p(q_j\setminus\omega_j, \mathbb{R}^2)} + \|\nabla u_j - \nabla a_j\|_{L^p(q_j\setminus\omega_j, \mathbb{R}^{2\times 2})} \leq c_p \|e(u)\|_{L^p(Q_j, \mathbb{R}^{2\times 2})}$, with a constant which depends only on the exponent p.

To pass to the estimate on $U^{\rm int}$ one uses a Whitney cover with pairs of open cubes q_j and Q_j such that the exterior ones have finite overlap and the interior ones form a cover, as done for example in proving the nonlinear Korn's inequality in [34, Theorem 3.1]. Following [33], if $\mathcal{H}^1(J_u \cap Q_j) \geq \eta r_j/8$ we define $P_j := x_j + (-r_j, r_j) \times (-r_j, \infty) \cap U$, otherwise $P_j = \emptyset$ and ω_j, a_j are obtained as above. Notice that $\mathcal{H}^1(P_j) \leq c_L r_j$ by the properties of Lipschitz functions and of the Whitney covering.

Then it suffices to apply the weighted Poincaré inequality, as done in [34, Theorem 3.1] and [33, Theorem 4.2]. By the properties of the covering, for neighboring pairs of squares $|q_j \cap q_i| \geq c r_i^2$, and if η is not too large this gives a bound on the difference of the two affine functions. One then defines $a^* \in C^\infty(U, \mathbb{R}^2)$ using a partition of unit subordinated to the cover $\{q_j\}$, and obtains

$$\int_{U^{\rm int}\setminus\cup_j P_j} (\varphi(x_1) - x_2)^p |D^2 a^*|^p(x) dx \leq c_{L,p} \|e(u)\|^p_{L^p(U, \mathbb{R}^2)}\,.$$

Since the cube $Q_0 = (-2,2)^2$ was not removed one has $a^* = a_0$ in q_0 and

application of the one-dimensional weighted Poincaré inequality to $a^*(x_1,\cdot)$ leads to the assertion, with $\omega := \cup_j (P_j \cup \omega_j)$ and $a := a_0$. Equivalently, in the last step one may use a Poincaré or Korn inequality on John domains, as done in [33, Theorem 4.2]. □

We remark that the nonoptimality of the exponent in [33, Theorem 4.2] is only consequence of the nonoptimal local estimate employed there (see [33, Theorem 3.1]).

3.3 Reflection

In this subsection we establish a technical result instrumental for the identification of the surface energy density in Section 4.3. To this aim, given $u \in SBD^p(\Omega)$ and a point $x_0 \in J_u$ we set

$$u_{x_0}(x) := \begin{cases} u^+(x_0) & \text{if } \langle x - x_0, \nu_{x_0}\rangle > 0, \\ u^-(x_0) & \text{if } \langle x - x_0, \nu_{x_0}\rangle < 0. \end{cases} \tag{3.6}$$

Lemma 3.3. *Let $p \in (1,\infty)$, $u \in SBD^p(\Omega)$, $\Omega \subset \mathbb{R}^2$ open. For $\mathcal{H}^1$-a.e. $x_0 \in J_u$ and any $\rho > 0$ sufficiently small there is $v_\rho \in SBD^p(B_{2\rho}(x_0)) \cap SBV^p(B_\rho(x_0),\mathbb{R}^2)$ such that:*

(i) $\lim_{\rho\to 0} \frac{1}{\rho}\mathcal{H}^1(J_{v_\rho} \setminus J_u \cap B_\rho(x_0)) = 0$;

(ii) $\lim_{\rho\to 0} \frac{1}{\rho}\int_{B_\rho(x_0)} |\nabla v_\rho|^p dx = 0$;

(iii) $\lim_{\rho\to 0} \frac{1}{\rho^2}\mathcal{L}^2(\{x \in B_\rho(x_0) : u \neq v_\rho\}) = 0$;

(iv) $\lim_{\rho\to 0} \frac{1}{\rho^2}\int_{B_\rho(x_0)} |v_\rho - u| dx = 0$;

(v) $\lim_{\rho\to 0} \frac{1}{\rho^{p+1}}\int_{B_\rho(x_0)} |v_\rho - u_{x_0}|^p dx = 0$;

(vi) $\lim_{\rho\to 0} \frac{1}{\rho}\int_{B_\rho(x_0)\cap J_{v_\rho}} |[v_\rho] - [u]| d\mathcal{H}^1 = 0$.

Proof. Since J_u is $(\mathcal{H}^1, 1)$ rectifiable, there exists a sequence $(\Gamma_i)_{i=1}^{\infty}$ of C^1 curves such that $\mathcal{H}^1(J_u \setminus \cup_{i=1}^{\infty}\Gamma_i) = 0$. For $\mathcal{H}^1$-a.e. $x_0 \in J_u$ we have

$$\lim_{\rho\to 0}\frac{1}{2\rho}\int_{J_u\cap B_\rho(x_0)}(|[u]|+1)d\mathcal{H}^1 = |[u](x_0)|+1,$$
$$\lim_{\rho\to 0}\frac{1}{2\rho}\int_{J_u\cap\Gamma\cap B_\rho(x_0)}(|[u]|+1)d\mathcal{H}^1 = |[u](x_0)|+1,$$

for one of the aforementioned curves Γ. Therefore

$$\lim_{\rho\to 0}\frac{1}{2\rho}\int_{(J_u\triangle\Gamma)\cap B_\rho(x_0)}(|[u]|+1)d\mathcal{H}^1 = 0 \tag{3.7}$$

and for ρ small Γ separates $B_{6\rho}(x_0)$ into two connected components. It is not restrictive to assume that $\Gamma \cap B_{6\rho}(x_0)$ is the graph of a function $h \in C^1(\mathbb{R})$. Moreover the following properties hold $\mathcal{H}^1$-a.e. $x_0 \in J_u$

$$\lim_{\rho\to 0}\frac{1}{\rho}\int_{B_\rho(x_0)}|e(u)|^p dx = 0, \tag{3.8}$$
$$\lim_{\rho\to 0}\frac{1}{2\rho}|Eu|(B_\rho(x_0)) = |[u]\odot\nu_u|(x_0), \tag{3.9}$$
$$\lim_{\rho\to 0}\frac{1}{\rho^2}\int_{B_\rho(x_0)\cap\{\pm(x_2-h(x_1))>0\}}|u-u^{\pm}(x_0)|dx = 0. \tag{3.10}$$

For simplicity we next assume that the point $x_0 = 0$ satisfies all the previous properties (3.7)-(3.10), with $h(0) = h'(0) = 0$. We also set $\tau_\rho := \|h\|_{L^\infty(B_{6\rho})}$ and note that $\tau_\rho/\rho \to 0$ as $\rho \to 0$. We now define the reflections of u with respect to the lines $\{x_2 = \pm\tau_\rho\}$, in the sense of [37, Lemma 1]. More precisely, define $\tilde{u}_\rho^+$ on the set $B_{2\rho}\cap\{x_2 < \tau_\rho\}$ by

$$\begin{cases}(\tilde{u}_\rho^+)_1(x_1,x_2) := -2u_1(x_1, 3\tau_\rho - 2x_2) + 3u_1(x_1, 2\tau_\rho - x_2)\\ (\tilde{u}_\rho^+)_2(x_1,x_2) := 4u_2(x_1, 3\tau_\rho - 2x_2) - 3u_2(x_1, 2\tau_\rho - x_2)\end{cases}$$

and by u otherwise in $B_{2\rho}$. Note that $\tilde{u}_\rho^+ \in SBD^p(B_{2\rho})$ and that

$$\lim_{\rho\to 0}\frac{1}{2\rho}\mathcal{H}^1(J_{\tilde{u}_\rho^+}\cap B_{2\rho}) = 0, \tag{3.11}$$
$$\|e(\tilde{u}_\rho^+)\|_{L^p(B_{2\rho},\mathbb{R}^{2\times 2})} \le c\|e(u)\|_{L^p(B_{6\rho},\mathbb{R}^{2\times 2})}, \tag{3.12}$$

for a universal constant c. Using a similar reflection we define $\tilde{u}_\rho^-$ in $B_{2\rho}\cap\{(x_1,x_2) : x_2 > -\tau_\rho\}$ and we set $\tilde{u}_\rho^- := u$ otherwise in $B_{2\rho}$.

By (3.7) and (3.11) for ρ small we have that $\tilde{u}_\rho^\pm$ satisfy the hypotheses of Proposition 3.2 on $B_{2\rho}$ with $s = 1/2$. Thus, there exist $w_\rho^\pm \in SBD^p(B_{2\rho})\cap$

$W^{1,p}(B_\rho,\mathbb{R}^2)$, for which properties (i)-(vii) hold true. Finally let us define $v_\rho \in SBD^p(B_{2\rho})$ by

$$v_\rho := \begin{cases} w_\rho^+ & \text{in } B_{2\rho}\cap\{x_2 > h(x_1)\},\\ w_\rho^- & \text{in } B_{2\rho}\cap\{x_2 < h(x_1)\}. \end{cases}$$

Since $w_\rho^\pm \in W^{1,p}(B_\rho,\mathbb{R}^2)$ we obtain $v_\rho \in SBV^p(B_\rho,\mathbb{R}^2)$ with

$$\begin{aligned} Dv_\rho \llcorner B_\rho =& \nabla w_\rho^+ \,\mathcal{L}^2 \llcorner B_\rho\cap\{x_2 > h(x_1)\} + (w_\rho^+ - w_\rho^-)\otimes \nu_\Gamma \mathcal{H}^1\llcorner \Gamma\cap B_\rho\\ &+ \nabla w_\rho^-\,\mathcal{L}^2\llcorner B_\rho\cap\{x_2 < h(x_1)\}. \end{aligned}$$

We next check that v_ρ satisfies the properties in the statement in the ball B_ρ. Property (i) comes straightforwardly from (3.7) and from the fact that $J_{v_\rho}\subset\Gamma$. Moreover (3.12), (3.3), and (3.8) yield

$$\lim_{\rho\to 0}\frac{1}{\rho}\int_{B_\rho}|e(w_\rho^\pm)|^p dx = 0. \tag{3.13}$$

As for property (iii), we observe that

$$\lim_{\rho\to 0}\frac{1}{\rho^2}\mathcal{L}^2(\{B_\rho :\, u\neq v_\rho\}) \le \lim_{\rho\to 0}(c\frac{\tau_\rho}{\rho}+\frac{c}{\rho}\mathcal{H}^1((J_u\setminus\Gamma)\cap B_{6\rho})) = 0,$$

where we have used Proposition 3.2 (i) and (3.7).

Let us now prove property (iv). By the definition of v_ρ and $\tilde u_\rho^\pm$ and by triangular inequality we obtain

$$\begin{aligned} &\frac{1}{\rho^2}\int_{B_\rho}|v_\rho-u|dx \le\\ &\quad\frac{1}{\rho^2}\int_{B_\rho\cap\{h(x_1)<x_2\}}|w_\rho^+-\tilde u_\rho^+|dx+\frac{1}{\rho^2}\int_{B_\rho\cap\{h(x_1)<x_2<\tau_\rho\}}|\tilde u_\rho^+-u|dx+\\ &\quad\frac{1}{\rho^2}\int_{B_\rho\cap\{x_2<h(x_1)\}}|w_\rho^--\tilde u_\rho^-|dx+\frac{1}{\rho^2}\int_{B_\rho\cap\{-\tau_\rho<x_2<h(x_1)\}}|\tilde u_\rho^--u|dx. \end{aligned}\tag{3.14}$$

By the definition of w_ρ^+ and Proposition 3.2 (vi) we can estimate

$$\frac{1}{\rho^2}\int_{B_\rho}|w_\rho^+-\tilde u_\rho^+|dx \le \frac{c}{\rho}|E\tilde u_\rho^+|(B_{2\rho}) \le \frac{c}{\rho}|Eu|(B_{6\rho}\setminus\Gamma).$$

By (3.7) and (3.8) we conclude that the first term of (3.14) tends to 0. Clearly, the same argument can be applied to the third term there. So, it

remains to treat the second term in (3.14), being the fourth one similar. By triangular inequality and a change of variable we infer

$$\frac{1}{\rho^2}\int_{B_\rho\cap\{h(x_1)<x_2<\tau_\rho\}}|\tilde{u}_\rho^+-u|dx\le$$
$$\frac{1}{\rho^2}\int_{B_\rho}|\tilde{u}_\rho^+-u^+(x_0)|dx+\frac{1}{\rho^2}\int_{B_\rho\cap\{h(x_1)<x_2\}}|u^+(x_0)-u|dx\le$$
$$\frac{c}{\rho^2}\int_{B_{6\rho}\cap\{h(x_1)<x_2\}}|u^+(x_0)-u|dx,$$

and the last term tends to 0 by (3.10), hence property (iv) follows.

Let us prove now property (v). By Korn's inequality and Poincaré's inequality in $W^{1,p}$, there exists an affine function $a_\rho(x):=d_\rho+\beta_\rho x$ such that

$$\frac{1}{\rho^{p+1}}\int_{B_\rho}|w_\rho^+-a_\rho|^p dx\le\frac{c}{\rho}\int_{B_\rho}|e(w_\rho^+)|^p dx. \tag{3.15}$$

We first claim that

$$\lim_{\rho\to0}d_\rho=u^+(x_0). \tag{3.16}$$

Let $\omega_\rho^+:=B_\rho\cap\{u=w_\rho^+\}\cap\{x_2>h(x_1)\}$. Since $|\omega_\rho^+|/\rho^2\to\pi/2$, and a_ρ is affine, by [21, Lemma 4.3] we obtain, for ρ small,

$$\|a_\rho-u^+(x_0)\|_{L^\infty(B_\rho^+,\mathbb{R}^2)}\le\frac{c}{\rho^2}\int_{\omega_\rho^+}|w_\rho^+-a_\rho|dx+\frac{c}{\rho^2}\int_{\omega_\rho^+}|u-u^+(x_0)|dx.$$

The right hand side above is infinitesimal by (3.15), (3.13) and (3.10), thus we conclude

$$\limsup_{\rho\to0}|d_\rho-u^+(x_0)|\le\lim_{\rho\to0}\|a_\rho-u^+(x_0)\|_{L^\infty(B_\rho^+,\mathbb{R}^2)}=0,$$

which proves (3.16).

Next we prove that

$$\lim_{\rho\to0}\rho|\beta_\rho|^p=0, \tag{3.17}$$
$$\lim_{\rho\to0}\rho^{\frac{1-p}{p}}|d_\rho-u^+(x_0)|=0. \tag{3.18}$$

To establish (3.17), we fix $\delta>0$ small and we consider $\hat\rho$ such that

$$\Big(\frac{1}{\rho}\int_{B_\rho}|e(w_\rho^+)|^p dx\Big)^{\frac{1}{p}}<\delta,\qquad\text{for }\rho\le\hat\rho, \tag{3.19}$$

note that this is possible by (3.13). For $\rho < \hat\rho$ we define $\rho_k := (2^k\rho) \wedge \hat\rho$ and we adopt the notation k in place of ρ_k for the subscriptions. As above, using [21, Lemma 4.3] and the triangular inequality we infer

$$\|a_k - a_{k+1}\|_{L^\infty(B^+_{\rho_k},\mathbb{R}^2)} \le \\ \frac{c}{{\rho_k}^2}\int_{\{u=w_k^+\}} |w_k^+ - a_k|dx + \frac{c}{\rho_{k+1}^2}\int_{\{u=w_{k+1}^+\}} |w_{k+1}^+ - a_{k+1}|dx \le c\delta\rho_k^{\frac{p-1}{p}},$$

where the last estimate follows by Hölder's inequality, (3.15), and (3.19). Therefore

$$|d_k - d_{k+1}| \le \|a_k - a_{k+1}\|_{L^\infty(B^+_{\rho_k},\mathbb{R}^2)} \le c\delta\rho_k^{\frac{p-1}{p}}, \tag{3.20}$$

and hence once more by [21, Lemma 4.3] and by the triangular inequality we conclude

$$|\beta_k - \beta_{k+1}| \le c\delta{\rho_k}^{-\frac{1}{p}}.$$

Collecting these estimates as k varies we obtain

$$\rho|\beta_\rho|^p \le \rho\Big(|\hat\beta| + \sum_{k=0}^{\hat k-1}|\beta_k - \beta_{k+1}|\Big)^p \le c\delta^p + c\rho|\hat\beta|^p,$$

where $\hat k$ is the first index such that $\rho_{\hat k} = \hat\rho$ and $\hat\beta := \beta_{\hat k} = \beta_{\hat\rho}$. This proves (3.17) as $\rho \to 0$ and $\delta \to 0$.

We next prove (3.18). Similarly to the previous estimate, summing (3.20) gives

$$|d_\rho - d_{\hat\rho}| \le c\delta\,\hat\rho^{(p-1)/p}$$

for all $0 < \rho < \hat\rho \le \rho_\delta$, with δ arbitrary and ρ_δ depending only on δ. Taking $\rho \to 0$ and using (3.16) yields

$$\hat\rho^{(1-p)/p}|u^+(x_0) - d_{\hat\rho}| \le c\delta$$

which, since δ was arbitrary, proves (3.18) and therefore (v).

At this point we turn to property (ii). Korn's inequality implies that

$$\|\nabla w_\rho^+\|_{L^p(B_\rho,\mathbb{R}^{2\times2})} \le \|\nabla w_\rho^+ - \beta_\rho\|_{L^p(B_\rho,\mathbb{R}^{2\times2})} + c\,\rho^{2/p}|\beta_\rho| \\ \le c\,\|e(w_\rho^+)\|_{L^p(B_\rho,\mathbb{R}^{2\times2})} + c\,\rho^{2/p}|\beta_\rho|,$$

where $c > 0$ is a universal constant. This, together with (3.13) and (3.17) and the corresponding estimates for w_ρ^-, implies property (ii).

We finally show property (vi). Note that by the trace theorem we have

$$\frac{1}{\rho}\int_{\Gamma\cap B_\rho}|v_\rho^\pm - u^\pm|d\mathcal{H}^1 \leq$$
$$\frac{c}{\rho^2}\int_{B_\rho}|v_\rho - u|dx + \frac{c}{\rho}|E(v_\rho - u)|(B_\rho \setminus \Gamma) \leq$$
$$\frac{c}{\rho^2}\int_{B_\rho}|v_\rho - u|dx + \frac{c}{\rho}\int_{B_\rho}|e(v_\rho)|dx + \frac{c}{\rho}\int_{B_\rho}|e(u)|dx + \frac{c}{\rho}\int_{J_u\setminus\Gamma}|[u]|d\mathcal{H}^1$$

and all terms in the last expression approach 0 respectively by (iv), (3.8), (3.13) and (3.7).

□

4 Integral representation

4.1 Preliminaries

In this Section we prove Theorem 1.1, along the lines of [9, Section 2.2].

Before starting we specify that property (ii) means that if $u_j, u \in SBD^p(\Omega)$ obey $u_j \to u$ in $L^1(\Omega,\mathbb{R}^2)$, then $F(u,A) \leq \liminf_{j\to\infty} F(u_j,A)$ for any open set A. Property (iii) means that if $u, v \in SBD^p(\Omega)$ obey $u = v$ $\mathcal{L}^2$-a.e. in A, then $F(u,A) = F(v,A)$. The functions f and g are defined in (4.1) and (4.2) below.

The family of balls contained in Ω is denoted by

$$\mathcal{A}^*(\Omega) := \{B_\varepsilon(x) : x \in \Omega, \varepsilon > 0\}\ .$$

Let $B \in \mathcal{A}^*(\Omega)$. We can identify any $u \in SBD^p(B)$ with its zero extension $u\chi_B \in SBD^p(\Omega)$, and correspondingly write $F(u,B)$ for $F(u\chi_B, B)$. By locality, for any other extension the value of the functional is the same.

For $B \in \mathcal{A}^*(\Omega)$ we define

$$m(u,B) := \inf\{F(w,B) : w \in SBD^p(B),\ w = u \text{ around } \partial B\}$$

where the condition $w = u$ around ∂B means that a ball $B' \subset\subset B$ exists, so that $w = u$ on $B \setminus B'$. For $\delta > 0$, $A \in \mathcal{A}(\Omega)$, we set

$$m^\delta(u,A) := \inf\{\sum_{i=1}^\infty m(u,B_i) : B_i \in \mathcal{A}^*, B_i \cap B_j = \emptyset, B_i \subset A,$$
$$\operatorname{diam}(B_i) < \delta,\ \mu(A \setminus \bigcup_{i=1}^\infty B_i) = 0\}\,,$$

where $\mu := \mathcal{L}^2 \llcorner \Omega + (1 + |[u]|)\mathcal{H}^1 \llcorner (J_u \cap \Omega)$.

Since $\delta \mapsto m^\delta(u, A)$ is decreasing, we can define

$$m^*(u, A) := \lim_{\delta \to 0} m^\delta(u, A).$$

Moreover, we set

$$f(x_0, u_0, \xi) := \limsup_{\varepsilon \to 0} \frac{m(u_0 + \xi(\cdot - x_0), B_\varepsilon(x_0))}{\mathcal{L}^2(B_\varepsilon)} \tag{4.1}$$

$$g(x_0, a, b, \nu) := \limsup_{\varepsilon \to 0} \frac{m(u_{x_0,a,b,\nu}, B_\varepsilon(x_0))}{2\varepsilon}, \tag{4.2}$$

where $u_{x_0,a,b,\nu}$ is defined as

$$u_{x_0,a,b,\nu}(x) := \begin{cases} a & \text{if } \langle x - x_0, \nu \rangle > 0, \\ b & \text{if } \langle x - x_0, \nu \rangle < 0. \end{cases}$$

Lemma 4.1. *For all $u \in SBD^p(\Omega)$ and $A \in \mathcal{A}(\Omega)$, $F(u, A) = m^*(u, A)$.*

Proof. By definition, $m(u, B) \le F(u, B)$ for any ball B. Since $F(u, \cdot)$ is a measure, we obtain $m^\delta(u, A) \le F(u, A)$ for any $\delta > 0$. Therefore $m^*(u, A) \le F(u, A)$.

To prove the converse inequality, let $\delta > 0$, pick countably many balls B_i^δ as in the definition of $m^\delta(u, A)$, such that

$$\sum_{i=1}^{\infty} m(u, B_i^\delta) < m^\delta(u, A) + \delta\,.$$

By the definition of m there are functions $v_i^\delta \in SBD^p(B_i^\delta)$ such that $v_i^\delta = u$ around ∂B_i^δ and $F(v_i^\delta, B_i^\delta) \le m(u, B_i^\delta) + \delta \mathcal{L}^2(B_i^\delta)$. We define

$$v^\delta := \sum_{i=1}^{\infty} v_i^\delta \chi_{B_i^\delta} + u \chi_{N_0^\delta}$$

where $N_0^\delta := \Omega \setminus \cup_i B_i^\delta$.

By the BD compactness theorem $v^\delta \in BD(\Omega)$ and by the SBD closure theorem (see also [26, Theorem 11.3]) we conclude that $v^\delta \in SBD^p(\Omega)$ and

$$Ev^\delta = \sum_{i=1}^{\infty} Ev_i^\delta \llcorner B_i^\delta + Eu \llcorner N_0^\delta\,,$$

with

$$|Ev^\delta| \llcorner N^\delta = 0, \quad \mu(N^\delta) = 0\,, \qquad F(v^\delta, N^\delta) = 0$$

where $N^\delta := A \cap N_0^\delta$. Further,

$$F(v^\delta, A) = \sum_{i=1}^{\infty} F(v_i^\delta, B_i^\delta) + F(v^\delta, N^\delta) \le m^\delta(u, A) + \delta + \delta \mathcal{L}^2(A)\,.$$

We claim that $v^\delta \to u$ in $L^1(\Omega, \mathbb{R}^2)$. Since $F(\cdot, A)$ is lower semicontinuous, this will imply

$$F(u, A) \le \liminf_{\delta \to 0} F(v^\delta, A) \le \liminf_{\delta \to 0} m^\delta(u, A) = m^*(u, A)\,.$$

To prove $v^\delta \to u$, we observe that by Poincaré's inequality, $\operatorname{diam} B_i^\delta \le \delta$, and $v^\delta = u$ on ∂B_i^δ we obtain

$$\|v^\delta - u\|_{L^1(B_i^\delta, \mathbb{R}^2)} \le c\delta |Ev^\delta - Eu|(B_i^\delta)\,.$$

Therefore

$$\begin{aligned}\|v^\delta - u\|_{L^1(\Omega, \mathbb{R}^2)} \le& \sum_i \|v^\delta - u\|_{L^1(B_i^\delta, \mathbb{R}^2)} \le c\delta(|Ev^\delta|(A) + |Eu|(A)) \\ \le& c\delta(F(v^\delta, A) + F(u, A))\,.\end{aligned}$$

Since $F(v^\delta, A)$ has a finite limit as $\delta \to 0$, this proves $v^\delta \to u$ in $L^1(\Omega, \mathbb{R}^2)$. □

Lemma 4.2. *For any ball $B_r(x_0) \subset \Omega$ and $\delta > 0$ sufficiently small we have*

(i) $\lim_{\delta \to 0} m(u, B_{r-\delta}(x_0)) = m(u, B_r(x_0))$;

(ii) $m(u, B_{r+\delta}(x_0))) \le m(u, B_r(x_0)) + \beta \int_{B_{r+\delta}(x_0) \setminus B_r(x_0)} (1 + |e(u)|^p) dx +$
$\beta \int_{J_u \cap B_{r+\delta}(x_0) \setminus B_r(x_0)} (1 + |[u]|) d\mathcal{H}^1$.

Proof. We drop x_0 from the notation. Choose $v_\delta \in SBD^p(B_{r-\delta})$ with $v_\delta = u$ around $\partial B_{r-\delta}$ and $F(v_\delta, B_{r-\delta}) \le m(u, B_{r-\delta}) + \delta$. We define

$$w_\delta(x) := \begin{cases} v_\delta(x) & \text{if } x \in B_{r-\delta}\,, \\ u(x) & \text{if } x \in \Omega \setminus B_{r-\delta}\,. \end{cases}$$

We have

$$\begin{aligned} m(u, B_r) \le& F(w_\delta, B_r) \le F(v_\delta, B_{r-\delta}) + F(w_\delta, B_r \setminus B_{r-\delta}) \\ \le& m(u, B_{r-\delta}) + \delta \\ &+ \beta \int_{B_r \setminus B_{r-\delta}} (|e(u)|^p + 1) dx + \beta \int_{J_u \cap B_r \setminus B_{r-\delta}} (1 + |[u]|) d\mathcal{H}^{n-1}\,. \end{aligned}$$

Since $(1+|e(u)|^p)\mathcal{L}^2 + (1+|[u]|)\mathcal{H}^1 \llcorner J_u$ is a bounded measure, we conclude that

$$m(u, B_r) \le \liminf_{\delta \to 0} m(u, B_{r-\delta})\,.$$

Conversely, for any $\varepsilon > 0$ there is $v_\varepsilon \in SBD^p(B_r)$ with $v_\varepsilon = u$ around ∂B_r and $F(v_\varepsilon, B_r) \le m(u, B_r) + \varepsilon$. For $\delta > 0$ sufficiently small one has $v_\varepsilon = u$ on $B_r \setminus B_{r-2\delta}$ and therefore $m(u, B_{r-\delta}) \le F(v_\varepsilon, B_{r-\delta}) \le m(u, B_r) + \varepsilon$. Taking first $\delta \to 0$ and then $\varepsilon \to 0$ concludes the proof of (i). The proof of (ii) is analogous. □

Lemma 4.3. *For μ-a.e. $x_0 \in \Omega$,*

$$\lim_{\varepsilon \to 0} \frac{F(u, B_\varepsilon(x_0))}{\mu(B_\varepsilon(x_0))} = \lim_{\varepsilon \to 0} \frac{m(u, B_\varepsilon(x_0))}{\mu(B_\varepsilon(x_0))}\,.$$

Proof. From $m(u, B_\varepsilon(x_0)) \le F(u, B_\varepsilon(x_0))$ one immediately obtains

$$\limsup_{\varepsilon \to 0} \frac{m(u, B_\varepsilon(x_0))}{\mu(B_\varepsilon(x_0))} \le \limsup_{\varepsilon \to 0} \frac{F(u, B_\varepsilon(x_0))}{\mu(B_\varepsilon(x_0))}$$

for any $x_0 \in \Omega$. To prove the converse inequality, we define for $t > 0$ the set

$$\begin{aligned} E_t := \{x \in \Omega : &\text{ there is } \varepsilon_h \to 0 \text{ such that} \\ &F(u, B_{\varepsilon_h}(x)) > m(u, B_{\varepsilon_h}(x)) + t\mu(B_{\varepsilon_h}(x)) \text{ for all } h\}\,. \end{aligned}$$

From this definition one immediately has

$$\liminf_{\varepsilon \to 0} \frac{F(u, B_\varepsilon(x_0))}{\mu(B_\varepsilon(x_0))} \le \liminf_{\varepsilon \to 0} \frac{m(u, B_\varepsilon(x_0))}{\mu(B_\varepsilon(x_0))} + t \quad \text{for all } x_0 \in \Omega \setminus E_t\,.$$

If we can prove that

$$\mu(E_t) = 0 \text{ for all } t > 0 \tag{4.3}$$

then, recalling that $\lim_{\varepsilon \to 0} \frac{F(u, B_\varepsilon(x_0))}{\mu(B_\varepsilon(x_0))}$ exists μ-almost everywhere, the proof is concluded.

It remains to prove (4.3) for an arbitrary $t > 0$. For $\delta > 0$ we define

$$\begin{aligned} X^\delta := \{B_\varepsilon(x) : \varepsilon < \delta,\ \overline{B}_\varepsilon(x) \subset \Omega,\ \mu(\partial B_\varepsilon(x)) = 0,\\ F(u, B_\varepsilon(x)) > m(u, B_\varepsilon(x)) + t\mu(B_\varepsilon(x))\} \end{aligned}$$

and

$$U^* := \bigcap_{\delta > 0} \{x : \exists \varepsilon > 0 \text{ s.t. } B_\varepsilon(x) \in X^\delta\}\,.$$

We first show that $E_t \subset U^*$. Let $x \in E_t$. Then for any $\delta > 0$ there is $\varepsilon \in (0, \delta)$ such that $F(u, B_\varepsilon(x)) > m(u, B_\varepsilon(x)) + t\mu(B_\varepsilon(x))$. By Lemma 4.2 the function $\varepsilon \to m(u, B_\varepsilon(x))$ is left-continuous; $F(u, B_\varepsilon(x))$ is left-continuous because $F(u, \cdot)$ is a measure, therefore the same inequality holds for all $\varepsilon' \in (\varepsilon'', \varepsilon)$. In particular, there is one which additionally obeys $\mu(\partial B_\varepsilon(x)) = 0$.

It remains to show that $\mu(U^*) = 0$. We fix a compact set $K \subset U^*$ and $0 < \delta < \eta$. Let $U^\eta := \bigcup\{B_\varepsilon(x) : B_\varepsilon(x) \in X^\eta\}$ and

$$Y^\delta := \{B_\varepsilon(x) : \varepsilon < \delta, \overline{B_\varepsilon(x)} \subset U^\eta \setminus K, \mu(\partial B_\varepsilon(x)) = 0\}\,.$$

By definition, X^δ is a fine cover of K and Y^δ of $U^\eta \setminus K$. Therefore there are countably many pairwise disjoint balls $B_i \in X^\delta$ and $\hat{B}_j \in Y^\delta$ and a set N with $\mu(N) = 0$ such that

$$U^\eta = \left(\bigcup_{i\in\mathbb{N}} B_i\right) \cup \left(\bigcup_{j\in\mathbb{N}} \hat{B}_j\right) \cup N\,.$$

Then

$$\begin{aligned} F(u, U^\eta) &= \sum_i F(u, B_i) + \sum_j F(u, \hat{B}_j) + F(u, N) \\ &\geq \sum_i (m(u, B_i) + t\mu(B_i)) + \sum_j m(u, \hat{B}_j) \\ &= \sum_i m(u, B_i) + \sum_j m(u, \hat{B}_j) + t\mu(\cup_i B_i) \\ &\geq m^\delta(u, U^\eta) + t\mu(K) \end{aligned}$$

where in the last step we used the definition of m^δ. For $\delta \to 0$, the definition of m^* and Lemma 4.1 give

$$F(u, U^\eta) \geq m^*(u, U^\eta) + t\mu(K) = F(u, U^\eta) + t\mu(K)\,.$$

Therefore $\mu(K) = 0$, and by the regularity of μ we conclude $\mu(U^*) = 0$.

□

4.2 Bounds on the volume term

In this subsection we identify the volume energy density in the integral representation for F to be the function f defined in (4.1). Throughout the whole subsection we consider a fixed map $u \in SBD^p(\Omega)$. Our first result shows that

the local volume energy density can be computed with a $W^{1,p}$-approximation to the blow-ups of u (see (4.7–4.8) below), in the sense that

$$\frac{dF(u,\cdot)}{d\mathcal{L}^2}(x_0) = \lim_{\varepsilon\to 0} \frac{m\big(w_\varepsilon, B_\varepsilon(x_0)\big)}{\mathcal{L}^2(B_\varepsilon)}. \tag{4.4}$$

We shall however not need (4.4), but only the apparently more complex version in (4.5)-(4.6). Taking a diagonal subsequence they imply (4.4).

Lemma 4.4. *For $\mathcal{L}^2$-almost any $x_0 \in \Omega$, any $\varepsilon > 0$, and any $s \in (0,1)$ there are functions $w_\varepsilon^s \in W^{1,p}(B_{s\varepsilon}(x_0);\mathbb{R}^2)$ which obey*

$$\frac{dF(u,\cdot)}{d\mathcal{L}^2}(x_0) \le \liminf_{s\to 1} \liminf_{\varepsilon\to 0} \frac{m\big(w_\varepsilon^s, B_{s\varepsilon}(x_0)\big)}{\mathcal{L}^2(B_{s\varepsilon})} \tag{4.5}$$

and

$$\limsup_{s\to 1} \limsup_{\varepsilon\to 0} \frac{m\big(w_\varepsilon^s, B_{s^2\varepsilon}(x_0)\big)}{\mathcal{L}^2(B_{s\varepsilon})} \le \frac{dF(u,\cdot)}{d\mathcal{L}^2}(x_0) \tag{4.6}$$

and which approximate the affine function $y \mapsto \nabla u(x_0)(y-x_0)+u(x_0)$ in the sense that

$$\lim_{\varepsilon\to 0} \frac{1}{\varepsilon^2} \int_{B_\varepsilon(x_0)} |e(w_\varepsilon^s) - e(u)(x_0)|^p dx = 0 \tag{4.7}$$

and

$$\lim_{\varepsilon\to 0} \frac{1}{\varepsilon^{2+p}} \int_{B_\varepsilon(x_0)} |w_\varepsilon^s(x) - u(x_0) - \nabla u(x_0)(x-x_0)|^p dx = 0\,. \tag{4.8}$$

We remark that the ball in (4.6) has radius $s^2\varepsilon$ instead of $s\varepsilon$. The estimate would also hold on $B_{s\varepsilon}$, the variant we chose is more convenient in the proof of Lemma 4.7.

Proof. Let $x_0 \in \Omega$ be such that

$$\lim_{\varepsilon\to 0} \frac{1}{\varepsilon^2} \int_{B_\varepsilon(x_0)} |e(u)(x) - e(u)(x_0)|^p dx = 0\,, \tag{4.9}$$

$$\lim_{\varepsilon\to 0} \frac{1}{\varepsilon^2} \int_{B_\varepsilon(x_0)\cap J_u} (1+|[u]|) d\mathcal{H}^1 = 0\,, \tag{4.10}$$

and

$$\lim_{\varepsilon\to 0} \frac{1}{\varepsilon^3} \int_{B_\varepsilon(x_0)} |u(x) - u(x_0) - \nabla u(x_0)(x-x_0)| dx = 0\,. \tag{4.11}$$

By [4, Th. 7.4], $\mathcal{L}^2$-almost every x_0 obeys (4.11), the other two are standard.

By (4.10), for sufficiently small ε one has $\mathcal{H}^1(J_u \cap B_\varepsilon(x_0)) \le \eta(1-s)\varepsilon/2$, where η is the constant from Theorem 2.1. By Proposition 3.2 applied to $u - u(x_0) - \nabla u(x_0)(\cdot - x_0)$ there is $\tilde{w}^s_\varepsilon \in SBD^p(B_\varepsilon(x_0)) \cap W^{1,p}(B_{s\varepsilon}(x_0);\mathbb{R}^2)$ with properties (i)-(vii) and we set $w^s_\varepsilon := \tilde{w}^s_\varepsilon + u(x_0) + \nabla u(x_0)(\cdot - x_0)$. In particular, (4.7) follows from (3.3) and (4.9), while (4.8) follows from Lemma 4.5 below applied to $\tilde{w}^s_\varepsilon$, estimating the right-hand side with (4.7), (vi), and (4.9)-(4.11).

We first prove (4.6). By the definition of m and the fact that $F(w^s_\varepsilon, \cdot)$ is a measure follows

$$m(w^s_\varepsilon, B_{s\varepsilon}(x_0)) \le F(w^s_\varepsilon, B_{s\varepsilon}(x_0)) \le F(w^s_\varepsilon, B_\varepsilon(x_0))\,.$$

Let $(B_i)_{i\in\mathbb{N}}$ be the balls from Proposition 3.2. For $M \in \mathbb{N}$ we define

$$w^{s,M}_\varepsilon := u + \chi_{\cup_{i=1}^M \overline{B}_i}(w^s_\varepsilon - u)\,.$$

Then $w^{s,M}_\varepsilon \in SBD^p(B_\varepsilon(x_0))$ and $w^{s,M}_\varepsilon \to w^s_\varepsilon$ in L^1 as $M \to \infty$. Further,

$$\begin{aligned} F(w^{s,M}_\varepsilon, B_\varepsilon(x_0)) \le& F(w^{s,M}_\varepsilon, B_\varepsilon(x_0) \setminus \cup_{i=1}^M \overline{B}_i) + \sum_{i=1}^M F(w^{s,M}_\varepsilon, \overline{B}_i) \\ \le& F(u, B_\varepsilon(x_0) \setminus \cup_{i=1}^M \overline{B}_i) + \beta \sum_{i=1}^M \int_{\overline{B}_i} (1 + |e(w^s_\varepsilon)|^p) dx \end{aligned}$$

since $w^{s,M}_\varepsilon = w^s_\varepsilon$ is a $W^{1,p}$ function on each $\overline{B}_i$. By monotonicity and lower semicontinuity of F we obtain

$$\begin{aligned} F(w^s_\varepsilon, B_\varepsilon(x_0)) \le& F(u, B_\varepsilon(x_0)) + \beta \sum_{i=1}^\infty \int_{\overline{B}_i} (1 + |e(w^s_\varepsilon)|^p) dx \\ \le& F(u, B_\varepsilon(x_0)) + c\mathcal{L}^2(\cup_i B_i)(1 + |e(u)|^p(x_0)) \\ &+ c \int_{B_\varepsilon(x_0)} (|e(w^s_\varepsilon) - e(u)(x_0)|^p) dx \end{aligned}$$

and, recalling Proposition 3.2 (i), conclude the proof of (4.6) by (4.7) and (4.9).

It remains to prove (4.5). Let $v_\varepsilon \in SBD^p(B_{s^2\varepsilon}(x_0))$ be such that $v_\varepsilon = w^s_\varepsilon$ around $\partial B_{s^2\varepsilon}(x_0)$ and $F(v_\varepsilon, B_{s^2\varepsilon}) \le m(w^s_\varepsilon, B_{s^2\varepsilon}(x_0)) + \varepsilon^3$. We define

$$\tilde{v}_\varepsilon(x) := \begin{cases} v_\varepsilon(x) & \text{if } x \in B_{s^2\varepsilon}(x_0) \\ w^s_\varepsilon(x) & \text{if } x \in B_\varepsilon(x_0) \setminus B_{s^2\varepsilon}(x_0)\,. \end{cases}$$

By definition of m and additivity of F we obtain

$$m(u, B_\varepsilon(x_0)) \leq F(\tilde{v}_\varepsilon, B_\varepsilon(x_0)) = F(\tilde{v}_\varepsilon, B_{s^2\varepsilon}(x_0)) + F(\tilde{v}_\varepsilon, B_\varepsilon(x_0) \setminus B_{s^2\varepsilon}(x_0))$$

where by locality of F and definition of v_ε

$$F(\tilde{v}_\varepsilon, B_{s^2\varepsilon}(x_0)) = F(v_\varepsilon, B_{s^2\varepsilon}(x_0)) \leq m(w^s_\varepsilon, B_{s^2\varepsilon}(x_0)) + \varepsilon^3$$

and, since $\tilde{v}_\varepsilon = w^s_\varepsilon$ outside $B_{s^2\varepsilon}(x_0)$ and $\mathcal{H}^1(J_{\tilde{v}_\varepsilon} \cap \partial B_{s^2\varepsilon}(x_0)) = 0$, recalling (3.3) we obtain

$$\begin{aligned}
F(\tilde{v}_\varepsilon, B_\varepsilon(x_0) \setminus B_{s^2\varepsilon}(x_0)) \leq & \beta \int_{B_\varepsilon(x_0)\setminus B_{s^2\varepsilon}(x_0)} (1 + |e(w^s_\varepsilon)|^p) dx \\
& + \beta \int_{J_u \cap B_\varepsilon(x_0)\setminus B_{s^2\varepsilon}(x_0)} (1 + |[u]|) d\mathcal{H}^1 \\
\leq & c\beta \mathcal{L}^2(B_\varepsilon)(1 - s^4)(1 + |e(u)|^p(x_0)) \\
& + c\beta \int_{B_\varepsilon(x_0)} |e(w^s_\varepsilon)(x) - e(u)(x_0)|^p dx \\
& + \beta \int_{J_u \cap B_\varepsilon(x_0)} (1 + |[u]|) d\mathcal{H}^1 \,.
\end{aligned}$$

Dividing by $\mathcal{L}^2(B_\varepsilon)$ and taking the limit $\varepsilon \to 0$ gives

$$\lim_{\varepsilon\to 0} \frac{m(u, B_\varepsilon(x_0))}{\mathcal{L}^2(B_\varepsilon)} \leq \liminf_{\varepsilon\to 0} \frac{m(w^s_\varepsilon, B_{s^2\varepsilon}(x_0))}{\mathcal{L}^2(B_\varepsilon)} + c\beta(1 - s^4)(1 + |e(u)|^p(x_0)) \,,$$

where we used (4.7) and (4.10). Recalling Lemma 4.3 we obtain

$$\frac{dF(u,\cdot)}{d\mathcal{L}^2}(x_0) = \lim_{\varepsilon\to 0} \frac{m(u, B_\varepsilon(x_0))}{\mathcal{L}^2(B_\varepsilon)} \leq \liminf_{s\to 1} \liminf_{\varepsilon\to 0} \frac{m(w^s_\varepsilon, B_{s^2\varepsilon}(x_0))}{\mathcal{L}^2(B_{s\varepsilon})} \,.$$

This concludes the proof of (4.5). □

The next Lemma is a reverse-Hölder estimate for functions with small strain, of the form $\|v\|_p \leq r\|e(v)\|_p + \|v\|_1 r^{-n/p'}$.

Lemma 4.5. *For any $p \geq 1$ there is $c > 0$ (depending on n and p) such that for any $v \in W^{1,p}(B_r; \mathbb{R}^n)$ one has*

$$\frac{1}{r^{n+p}} \int_{B_r} |v|^p dx \leq c \frac{1}{r^n} \int_{B_r} |e(v)|^p dx + c \left(\frac{1}{r^{n+1}} \int_{B_r} |v| dx \right)^p \,.$$

Proof. By scaling it suffices to consider $r = 1$. By Korn's inequality there is an affine function a such that

$$\int_{B_1} |v - a|^p dx \le c \int_{B_1} |e(v)|^p dx \,.$$

Since a is affine,

$$\int_{B_1} |a|^p dx \le c \left(\int_{B_1} |a| dx \right)^p \le c \left(\int_{B_1} |v| dx \right)^p + c \int_{B_1} |v - a|^p dx \,.$$

A triangular inequality concludes the proof. □

Lemma 4.6. *For $\mathcal{L}^2$-a.e. $x_0 \in \Omega$,*

$$\frac{dF(u,\cdot)}{d\mathcal{L}^2}(x_0) \le f(x_0, u(x_0), \nabla u(x_0))$$

where f was defined in (4.1).

Proof. Let x_0, w^s_ε be as in Lemma 4.4, for $s \in (0,1)$. We choose $v^s_\varepsilon \in SBD^p(B_{s^2\varepsilon}(x_0))$ such that $v^s_\varepsilon(x) = u(x_0) + \nabla u(x_0)(x - x_0)$ around $\partial B_{s^2\varepsilon}(x_0)$ and $F(v^s_\varepsilon, B_{s^2\varepsilon}(x_0)) \le m(u(x_0) + \nabla u(x_0)(\cdot - x_0), B_{s^2\varepsilon}(x_0)) + \varepsilon^3$. We extend it to $\mathbb{R}^2$ setting it equal to $u(x_0) + \nabla u(x_0)(\cdot - x_0)$ outside $B_{s^2\varepsilon}(x_0)$ and choose $\varphi \in C^\infty_c(B_{s\varepsilon}(x_0))$ with $\varphi = 1$ on $B_{s^2\varepsilon}(x_0)$ and $\|D\varphi\|_\infty \le c/(s(1-s)\varepsilon)$. We define

$$z^s_\varepsilon := \varphi v^s_\varepsilon + (1 - \varphi) w^s_\varepsilon \,.$$

We remark that $z^s_\varepsilon = v^s_\varepsilon$ on $B_{s^2\varepsilon}(x_0)$ and $z^s_\varepsilon \in W^{1,p}(B_{s\varepsilon}(x_0) \setminus B_{s^2\varepsilon}(x_0); \mathbb{R}^2)$. Then

$$\begin{aligned} m(w^s_\varepsilon, B_{s\varepsilon}(x_0)) \le & F(z^s_\varepsilon, B_{s\varepsilon}(x_0)) \le F(v^s_\varepsilon, B_{s^2\varepsilon}(x_0)) + F(z^s_\varepsilon, B_{s\varepsilon}(x_0) \setminus B_{s^2\varepsilon}(x_0)) \\ \le & m(u(x_0) + \nabla u(x_0)(\cdot - x_0), B_{s^2\varepsilon}(x_0)) + \varepsilon^3 \\ & + \beta \int_{B_{s\varepsilon}(x_0) \setminus B_{s^2\varepsilon}(x_0)} (1 + |e(z^s_\varepsilon)|^p) dx \,. \end{aligned}$$

In order to estimate the error term, we observe that in $B_{s\varepsilon}(x_0) \setminus B_{s^2\varepsilon}(x_0)$ one has

$$\nabla z^s_\varepsilon - \nabla u(x_0) = (u(x_0) + \nabla u(x_0)(\cdot - x_0) - w^s_\varepsilon) \nabla \varphi + (1 - \varphi)(\nabla u(x_0) - \nabla w^s_\varepsilon)$$

which implies

$$\begin{aligned} \int_{B_{s\varepsilon}(x_0) \setminus B_{s^2\varepsilon}(x_0)} (1 + |e(z^s_\varepsilon)|^p) dx \le & c(1-s)\mathcal{L}^2(B_{s\varepsilon})(1 + |e(u)|^p(x_0)) \\ & + c \int_{B_{s\varepsilon}(x_0)} |e(u)(x_0) - e(w^s_\varepsilon)|^p dx \\ & + c \int_{B_{s\varepsilon}(x_0)} \frac{|u(x_0) + \nabla u(x_0)(\cdot - x_0) - w^s_\varepsilon|^p}{\varepsilon^p s^p (1-s)^p} dx. \end{aligned}$$

Therefore

$$\limsup_{\varepsilon\to 0}\frac{F(z^s_\varepsilon, B_{s\varepsilon}(x_0)\setminus B_{s^2\varepsilon}(x_0))}{\mathcal{L}^2(B_{s\varepsilon})}\le c(1-s)(1+|e(u)|^p(x_0))$$

and

$$\begin{aligned}\limsup_{\varepsilon\to 0}\frac{m(w^s_\varepsilon, B_{s\varepsilon}(x_0))}{\mathcal{L}^2(B_{s\varepsilon})}\le&\limsup_{\varepsilon\to 0}\frac{m(u(x_0)+\nabla u(x_0)(\cdot-x_0), B_{s^2\varepsilon}(x_0))}{\mathcal{L}^2(B_{s\varepsilon})}\\&+c(1-s)(1+|e(u)|^p(x_0))\\=&s^2 f(x_0,u_0,\nabla u(x_0))+c(1-s)(1+|e(u)|^p(x_0))\,.\end{aligned}$$

Since s was arbitrary, this concludes the proof. □

Lemma 4.7. *For $\mathcal{L}^2$-a.e. $x_0\in\Omega$,*

$$f(x_0,u(x_0),\nabla u(x_0))\le\frac{dF(u,\cdot)}{d\mathcal{L}^2}(x_0)$$

where f was defined in (4.1).

Proof. We choose x_0 and w^s_ε as in Lemma 4.4, for $s\in(0,1)$. We let $v^s_\varepsilon\in SBD^p(B_{s^2\varepsilon}(x_0))$ be such that $v^s_\varepsilon=w^s_\varepsilon$ around $\partial B_{s^2\varepsilon}(x_0)$ and $F(v^s_\varepsilon,B_{s^2\varepsilon}(x_0))\le m(w^s_\varepsilon,B_{s^2\varepsilon}(x_0))+\varepsilon^3$, and extend it to $B_{s\varepsilon}(x_0)$ setting it equal to w^s_ε outside $B_{s^2\varepsilon}(x_0)$. We choose $\varphi\in C^\infty_c(B_{s\varepsilon}(x_0))$ with $\varphi=1$ on $B_{s^2\varepsilon}(x_0)$ and $\|D\varphi\|_\infty\le c/(s(1-s)\varepsilon)$ and define

$$z^s_\varepsilon:=\varphi v^s_\varepsilon+(1-\varphi)(u(x_0)+\nabla u(x_0)(x-x_0))\,.$$

Then

$$\begin{aligned}m(u(x_0)+\nabla u(x_0)(\cdot-x_0),&B_{s\varepsilon}(x_0))\le F(z^s_\varepsilon,B_{s\varepsilon}(x_0))\\=&F(v^s_\varepsilon,B_{s^2\varepsilon}(x_0))+F(z^s_\varepsilon,B_{s\varepsilon}(x_0)\setminus B_{s^2\varepsilon}(x_0))\\\le&m(w^s_\varepsilon,B_{s^2\varepsilon}(x_0))+\varepsilon^3+F(z^s_\varepsilon,B_{s\varepsilon}(x_0)\setminus B_{s^2\varepsilon}(x_0))\,.\end{aligned}$$

In order to estimate the error term, we observe that in $B_{s\varepsilon}(x_0)\setminus B_{s^2\varepsilon}(x_0)$ one has

$$\nabla z^s_\varepsilon-\nabla u(x_0)=-(u(x_0)+\nabla u(x_0)(\cdot-x_0)-w^s_\varepsilon)\nabla\varphi+\varphi(\nabla w^s_\varepsilon-\nabla u(x_0))$$

which leads as in the proof of Lemma 4.6 to

$$\limsup_{\varepsilon\to 0}\frac{F(z^s_\varepsilon,B_{s\varepsilon}(x_0)\setminus B_{s^2\varepsilon}(x_0))}{\mathcal{L}^2(B_{s\varepsilon})}\le c(1-s)(1+|e(u)|^p(x_0))\,.$$

We conclude that for any $s \in (0,1)$

$$\limsup_{\varepsilon\to 0} \frac{m(u(x_0)+\nabla u(x_0)(\cdot - x_0), B_{s\varepsilon}(x_0))}{\mathcal{L}^2(B_{s\varepsilon})}$$
$$\leq \limsup_{\varepsilon\to 0} \frac{m(w_\varepsilon^s, B_{s^2\varepsilon}(x_0))}{\mathcal{L}^2(B_{s\varepsilon})} + c(1-s)(1+|e(u)|^p(x_0))\,.$$

Since s was arbitrary, this concludes the proof. □

4.3 Bounds on the surface term

In the current subsection we identify the function g in (4.2) to be the surface energy density in the integral representation of F. As above, we work with a fixed map $u \in SBD^p(\Omega)$.

We first prove a technical result.

Lemma 4.8. *For $\mathcal{H}^1$-a.e. $x_0 \in J_u$ the functions $v_{2\varepsilon} \in SBV^p(B_{2\varepsilon}(x_0), \mathbb{R}^2)$ introduced in Lemma 3.3 satisfy for all $t \in (0,2)$*

$$\frac{dF(u,\cdot)}{d\mathcal{H}^1 \llcorner J_u}(x_0) = \lim_{\varepsilon\to 0} \frac{m(v_{2\varepsilon}, B_{t\varepsilon}(x_0))}{2t\varepsilon}. \tag{4.12}$$

Proof. It suffices to consider points x_0 such that the conclusions of Lemmata 3.3 and 4.3 hold true, the Radon-Nikodym derivative $\frac{dF(u,\cdot)}{d\mathcal{H}^1 \llcorner J_u}(x_0)$ exists finite,

$$\lim_{\varepsilon\to 0} \frac{\mu(B_\varepsilon(x_0))}{2\varepsilon} = 1 + |[u](x_0)|, \tag{4.13}$$

and

$$\lim_{\varepsilon\to 0} \Big(\frac{1}{\varepsilon} \int_{B_\varepsilon(x_0)} |e(u)|^p dx + \frac{1}{\varepsilon^2} \int_{B_\varepsilon(x_0)} |u(x) - u_{x_0}| dx \Big) = 0, \tag{4.14}$$

where u_{x_0} is the piecewise constant function defined in (3.6). In view of all these choices and thanks to Lemma 4.3 we may conclude that

$$\frac{dF(u,\cdot)}{d\mathcal{H}^1 \llcorner J_u}(x_0) = \lim_{\varepsilon\to 0} \frac{F(u, B_\varepsilon(x_0))}{2\varepsilon} = \lim_{\varepsilon\to 0} \frac{m(u, B_\varepsilon(x_0))}{2\varepsilon}. \tag{4.15}$$

For $\varepsilon > 0$ small enough the function $v_{2\varepsilon}$ introduced in Lemma 3.3 belongs to $SBD^p(B_{4\varepsilon}(x_0)) \cap SBV^p(B_{2\varepsilon}(x_0), \mathbb{R}^2)$ and it satisfies properties (i)-(vi). We set $w_\varepsilon := v_{2\varepsilon}$, we are left with proving that for all $t \in (0,2)$

$$\frac{dF(u,\cdot)}{d\mathcal{H}^1 \llcorner J_u}(x_0) \geq \limsup_{\varepsilon\to 0} \frac{m(w_\varepsilon, B_{t\varepsilon}(x_0))}{2t\varepsilon}, \tag{4.16}$$

$$\frac{dF(u,\cdot)}{d\mathcal{H}^1 \llcorner J_u}(x_0) \le \liminf_{\varepsilon\to 0} \frac{m(w_\varepsilon, B_{t\varepsilon}(x_0))}{2t\varepsilon}. \tag{4.17}$$

For the sake of notational simplicity we shall prove inequalities (4.16) and (4.17) only for $t = 1$.

We start off with (4.16). Let $(\varepsilon_j)_j$ be a sequence such that

$$\lim_{j\to\infty} \frac{m(w_{\varepsilon_j}, B_{\varepsilon_j}(x_0))}{2\varepsilon_j} = \limsup_{\varepsilon\to 0} \frac{m(w_\varepsilon, B_\varepsilon(x_0))}{2\varepsilon}. \tag{4.18}$$

Items (iii) and (iv) in Lemma 3.3 and the Coarea formula yield for a subsequence not relabeled for convenience that for $\mathcal{L}^1$-a.e. $s \in (0,1)$

$$\lim_j \frac{1}{\varepsilon_j} \int_{\partial B_{s\varepsilon_j}(x_0)\cap\{u\neq w_{\varepsilon_j}\}} \big(1 + |u - w_{\varepsilon_j}|\big) d\mathcal{H}^1 = 0, \tag{4.19}$$

$$\mu\big(\partial B_{s\varepsilon_j}(x_0)\big) = \mathcal{H}^1\big(\partial B_{s\varepsilon_j}(x_0) \cap J_{w_{\varepsilon_j}}\big) = 0. \tag{4.20}$$

We choose $z_j \in SBD^p(B_{s\varepsilon_j}(x_0))$ such that $z_j = u$ around $\partial B_{s\varepsilon_j}(x_0)$ and

$$F(z_j, B_{s\varepsilon_j}(x_0)) \le m(u, B_{s\varepsilon_j}(x_0)) + \varepsilon_j^2,$$

and define

$$\zeta_j := \begin{cases} z_j & B_{s\varepsilon_j}(x_0) \\ w_{\varepsilon_j} & B_{\varepsilon_j}(x_0) \setminus \overline{B_{s\varepsilon_j}(x_0)}. \end{cases}$$

The definition of z_j, the growth conditions in (1.1), and the locality of F yield

$$\begin{aligned} m(w_{\varepsilon_j}, B_{\varepsilon_j}(x_0)) &\le F(\zeta_j, B_{\varepsilon_j}(x_0)) \\ &\le F(z_j, B_{s\varepsilon_j}(x_0)) + \underbrace{\beta \int_{B_{\varepsilon_j}(x_0)\setminus B_{s\varepsilon_j}(x_0)} (1 + |e(w_{\varepsilon_j})|^p)\, dx}_{=:I_j^{(1)}} \\ &+ \underbrace{\beta \int_{\partial B_{s\varepsilon_j}(x_0)\cap\{u\neq w_{\varepsilon_j}\}} (1 + |u - w_{\varepsilon_j}|) d\mathcal{H}^1}_{=:I_j^{(2)}} + \underbrace{\beta \int_{(B_{\varepsilon_j}(x_0)\setminus\overline{B_{s\varepsilon_j}(x_0)})\cap J_{w_{\varepsilon_j}}} (1 + |[w_{\varepsilon_j}]|) d\mathcal{H}^1}_{=:I_j^{(3)}} \\ &\le m(u, B_{s\varepsilon_j}(x_0)) + \varepsilon_j^2 + I_j^{(1)} + I_j^{(2)} + I_j^{(3)}. \end{aligned}$$

We note that $I_j^{(1)}$ and $I_j^{(2)}$ are $o(\varepsilon_j)$ as $j \to \infty$ thanks to Lemma 3.3 (ii) and (4.19), respectively. Instead, employing Lemma 3.3 (vi) and (4.13) to bound

$I_j^{(3)}$ we infer that

$$\limsup_{j\to\infty}\frac{I_j^{(3)}}{2\varepsilon_j}\le\limsup_{j\to\infty}\frac{\beta}{2\varepsilon_j}\int_{(B_{\varepsilon_j}(x_0)\setminus B_{s\varepsilon_j}(x_0)\cap J_u}(1+|[u]|)d\mathcal{H}^1$$
$$=\beta\limsup_{j\to\infty}\frac{\mu\big(B_{\varepsilon_j}(x_0)\setminus B_{s\varepsilon_j}(x_0)\cap J_u\big)}{2\varepsilon_j}=(1-s)\beta(1+|[u](x_0)|). \quad (4.21)$$

Therefore, by (4.15) we conclude

$$\lim_{j\to\infty}\frac{m(w_{\varepsilon_j},B_{\varepsilon_j}(x_0))}{2\varepsilon_j}\le\liminf_{j\to\infty}\frac{m(u,B_{s\varepsilon_j}(x_0))}{2\varepsilon_j}+(1-s)\beta(1+|[u](x_0)|)$$
$$=s\frac{dF(u,\cdot)}{d\mathcal{H}^1\llcorner J_u}(x_0)+(1-s)\beta(1+|[u](x_0)|).$$

Estimate (4.16) follows at once by (4.18) and by letting $s\uparrow 1$ in the last inequality.

Let now $(\varepsilon_j)_j$ be a sequence such that

$$\lim_{j\to\infty}\frac{m(w_{\varepsilon_j},B_{\varepsilon_j}(x_0))}{2\varepsilon_j}=\liminf_{\varepsilon\to 0}\frac{m(w_\varepsilon,B_\varepsilon(x_0))}{2\varepsilon}. \quad (4.22)$$

Let $\lambda\in(1,2)$, arguing as for (4.19) and (4.20), up to a subsequence depending on λ and not relabeled for convenience we may assume that for $\mathcal{L}^1$-a.e. $s\in(0,1)$

$$\lim_{j\to\infty}\frac{1}{\varepsilon_j}\int_{\partial B_{s\lambda\varepsilon_j}(x_0)\cap\{u\neq w_{\varepsilon_j}\}}\big(1+|u-w_{\varepsilon_j}|\big)d\mathcal{H}^1=0, \quad (4.23)$$

and

$$\mu\big(\partial B_{s\lambda\varepsilon_j}(x_0)\big)=\mathcal{H}^1\big(\partial B_{s\lambda\varepsilon_j}(x_0)\cap J_{w_{\varepsilon_j}}\big)=0. \quad (4.24)$$

Given $z_j\in SBD^p(B_{s\lambda\varepsilon_j}(x_0))$ with $z_j=w_{\varepsilon_j}$ around $\partial B_{s\lambda\varepsilon_j}(x_0)$ and such that

$$F(z_j,B_{s\lambda\varepsilon_j}(x_0))\le m(w_{\varepsilon_j},B_{s\lambda\varepsilon_j}(x_0))+\varepsilon_j^2,$$

define

$$\zeta_j:=\begin{cases}z_j & B_{s\lambda\varepsilon_j}(x_0)\\ u & B_{\lambda\varepsilon_j}(x_0)\setminus\overline{B_{s\lambda\varepsilon_j}(x_0)}.\end{cases}$$

Using ζ_j as a test field for $m(u,B_{\lambda\varepsilon_j}(x_0))$, by the locality of F and its growth

conditions in (1.1)

$$m(u, B_{\lambda\varepsilon_j}(x_0)) \le F(\zeta_j, B_{\lambda\varepsilon_j}(x_0)) \le m(w_{\varepsilon_j}, B_{s\lambda\varepsilon_j}(x_0)) + \varepsilon_j^2$$
$$+ \underbrace{\beta \int_{B_{\lambda\varepsilon_j}(x_0)} (1 + |e(u)|^p)\, dx}_{I_j^{(4)}} + \underbrace{\beta \int_{\partial B_{s\lambda\varepsilon_j}(x_0) \cap \{u \neq w_{\varepsilon_j}\}} (1 + |u - w_{\varepsilon_j}|) d\mathcal{H}^1}_{I_j^{(5)}}$$
$$+ \underbrace{\beta \int_{(B_{\lambda\varepsilon_j}(x_0) \setminus \overline{B_{s\lambda\varepsilon_j}(x_0)}) \cap J_u} (1 + |[u]|) d\mathcal{H}^1}_{I_j^{(6)}}.$$

The terms $I_j^{(4)}$ and $I_j^{(5)}$ are $o(\varepsilon_j)$ by (4.14) and (4.23), respectively. The term $I_j^{(6)}$ can be estimated thanks to (4.13). Hence, we get by (4.15)

$$\frac{dF(u,\cdot)}{d\mathcal{H}^1 \llcorner J_u}(x_0) = \limsup_{j\to\infty} \frac{m(u, B_{\lambda\varepsilon_j}(x_0))}{2\lambda\varepsilon_j} \le \limsup_{j\to\infty} \frac{m(w_{\varepsilon_j}, B_{s\lambda\varepsilon_j}(x_0))}{2\lambda\varepsilon_j}. \quad (4.25)$$

Next, by choosing $s \in (0,1)$ for which (4.23) and (4.24) hold and $s\lambda > 1$, we may use Lemma 4.2(ii) to infer

$$m(w_{\varepsilon_j}, B_{s\lambda\varepsilon_j}(x_0)) \le m(w_{\varepsilon_j}, B_{\varepsilon_j}(x_0)) + \beta \int_{B_{s\lambda\varepsilon_j}(x_0) \setminus B_{\varepsilon_j}(x_0)} (1 + |e(w_{\varepsilon_j})|^p) dx$$
$$+ \beta \int_{(B_{s\lambda\varepsilon_j}(x_0) \setminus B_{\varepsilon_j}(x_0)) \cap J_{w_{\varepsilon_j}}} (1 + |[w_{\varepsilon_j}]|) d\mathcal{H}^1. \quad (4.26)$$

Clearly, the first integral is $o(\varepsilon_j)$ by Lemma 3.3 (ii), while the other one can be dealt with as $I_j^{(3)}$ in (4.21). Thus, (4.25) and (4.26) give

$$\frac{dF(u,\cdot)}{d\mathcal{H}^1 \llcorner J_u}(x_0) \le \frac{1}{\lambda} \lim_{j\to\infty} \frac{m(w_{\varepsilon_j}, B_{\varepsilon_j}(x_0))}{2\varepsilon_j} + (s\lambda - 1)\beta(1 + |[u](x_0)|).$$

In conclusion, by taking into account (4.22), we deduce (4.17) by taking first the limit as $s \uparrow 1$, for $s \in (0,1)$ chosen as explained above, and then as $\lambda \downarrow 1$ in the latter inequality. □

We are now ready to show that the function g in (4.2) is the surface energy density of F. This task shall be accomplished by proving two inequalities.

Lemma 4.9. *For $\mathcal{H}^1$-a.e. $x_0 \in J_u$,*

$$\frac{dF(u,\cdot)}{d\mathcal{H}^1 \llcorner J_u}(x_0) \le g(x_0, u^+(x_0), u^-(x_0), \nu_u(x_0))$$

where g was defined in (4.2).

Proof. We consider the same x_0 as in Lemma 4.8. In view of (4.12) and the definition of g in (4.2) it suffices to show that

$$\lim_{\varepsilon\to 0}\frac{m(w_\varepsilon, B_\varepsilon(x_0))}{2\varepsilon} \le \limsup_{\varepsilon\to 0}\frac{m(u_{x_0}, B_\varepsilon(x_0))}{2\varepsilon}, \tag{4.27}$$

where w_ε is the function introduced in Lemma 4.8. To prove such a claim consider any sequence $(\varepsilon_j)_j$, we have that for $\mathcal{L}^1$-a.e. $s\in(0,1)$

$$\mu\big(\partial B_{s\varepsilon_j}(x_0)\big) = \mathcal{H}^1\big(\partial B_{s\varepsilon_j}(x_0)\cap J_{w_j}\big) = 0, \tag{4.28}$$

where we have set $w_j := w_{\varepsilon_j}$.

Fix $s\in(0,1)$ as above and a test field $z_j\in SBD^p(B_{s\varepsilon_j}(x_0))$ with $z_j = u_{x_0}$ on $\partial B_{s\varepsilon_j}(x_0)$ such that

$$F(z_j, B_{s\varepsilon_j}(x_0)) \le m(u_{x_0}, B_{s\varepsilon_j}(x_0)) + \varepsilon_j^2.$$

Consider a cut-off function $\varphi\in C_c^\infty(B_{\varepsilon_j}(x_0),[0,1])$ such that $\varphi\equiv 1$ on $B_{s\varepsilon_j}(x_0)$ and $\|\nabla\varphi\|_{L^\infty}\le\frac{2}{(1-s)\varepsilon_j}$. Define $\zeta_j := \varphi\, z_j + (1-\varphi)w_j$, with the convention that z_j is extended equal to u_{x_0} outside $B_{s\varepsilon_j}(x_0)$. Therefore, by using ζ_j as a test field for $m(w_j, B_{\varepsilon_j}(x_0))$ we infer from the growth condition in (1.1) and the locality of F

$$\begin{aligned}
&m(w_j, B_{\varepsilon_j}(x_0)) \le F(\zeta_j, B_{\varepsilon_j}(x_0)) \le F(z_j, B_{s\varepsilon_j}(x_0))\\
&+\underbrace{C\int_{B_{\varepsilon_j}(x_0)\setminus B_{s\varepsilon_j}(x_0)}(1+|e(w_j)|^p)\,dx}_{=:I_j^{(7)}} + \underbrace{\frac{C}{((1-s)\varepsilon_j)^p}\int_{B_{\varepsilon_j}(x_0)\setminus B_{s\varepsilon_j}(x_0)}|w_j-u_{x_0}|^p dx}_{=:I_j^{(8)}}\\
&+\underbrace{C\,\mathcal{H}^1\big((B_{\varepsilon_j}(x_0)\setminus B_{s\varepsilon_j}(x_0))\cap J_{\zeta_j}\big)}_{=:I_j^{(9)}} + \underbrace{C\int_{(B_{\varepsilon_j}(x_0)\setminus B_{s\varepsilon_j}(x_0))\cap J_{\zeta_j}}|[\zeta_j]|d\mathcal{H}^1}_{=:I_j^{(10)}}\\
&\le m(u_{x_0}, B_{s\varepsilon_j}(x_0)) + \varepsilon_j^2 + I_j^{(7)} + I_j^{(8)} + I_j^{(9)} + I_j^{(10)},
\end{aligned} \tag{4.29}$$

with $C = C(\beta,p) > 0$.

By taking into account Lemma 3.3 (ii) and (v) we deduce that $I_j^{(7)}+I_j^{(8)} = o(\varepsilon_j)$ as $j\to\infty$. Moreover, as

$$\mathcal{H}^1((B_{\varepsilon_j}(x_0)\setminus B_{s\varepsilon_j}(x_0))\cap J_{\zeta_j}\setminus(J_{u_{x_0}}\cup J_{w_j})) = 0,$$

item (i) in Lemma 3.3 together with (4.13) give

$$\limsup_{j\to\infty}\frac{I_j^{(9)}}{2\varepsilon_j} \le C(1-s)(1+|[u](x_0)|).$$

Furthermore, for $\mathcal{H}^1$-a.e. $x \in J_{\zeta_j} \cap (B_{\varepsilon_j}(x_0) \setminus \overline{B_{s\varepsilon_j}(x_0)})$ it holds

$$|[\zeta_j]| \le |[u_{x_0}]|\chi_{J_{u_{x_0}} \cap J_{\zeta_j}} + |[w_j]|\chi_{J_{w_j} \cap J_{\zeta_j}} \le 2|[u_{x_0}]|\chi_{J_{\zeta_j}} + |[w_j] - [u_{x_0}]|\chi_{J_{w_j}}.$$

In turn the latter inequality implies by (4.13) and (4.28)

$$\begin{aligned}\limsup_{j\to\infty} \frac{I_j^{(10)}}{2\varepsilon_j} &\le C(1-s)|[u](x_0)| \\ &\quad + C \limsup_{j\to\infty} \frac{1}{2\varepsilon_j} \int_{(B_{\varepsilon_j}(x_0)\setminus B_{s\varepsilon_j}(x_0))\cap J_{\zeta_j}} (|[w_j] - [u](x_0)|)\, d\mathcal{H}^1 \\ &\le C(1-s)|[u](x_0)|,\end{aligned}$$

thanks to item (vi) in Lemma 3.3.

Finally, we obtain from (4.29)

$$\begin{aligned}\liminf_{j\to\infty} \frac{m(w_j, B_{\varepsilon_j}(x_0))}{2\varepsilon_j} &\le s \limsup_{j\to\infty} \frac{m(u_{x_0}, B_{s\varepsilon_j}(x_0))}{2s\varepsilon_j} + C(1-s)(1+|[u](x_0)|) \\ &\le s \limsup_{\varepsilon\to 0} \frac{m(u_{x_0}, B_{\varepsilon}(x_0))}{2\varepsilon} + C(1-s)(1+|[u](x_0)|),\end{aligned}$$

and the claim in (4.27) follows at once by letting $s \to 1$ in the inequality above. □

The reverse inequality is established arguing in an analogous fashion, therefore we provide a more concise proof.

Lemma 4.10. *For $\mathcal{H}^1$-a.e. $x_0 \in J_u$,*

$$\frac{dF(u,\cdot)}{d\mathcal{H}^1 \llcorner J_u}(x_0) \ge g(x_0, u^+(x_0), u^-(x_0), \nu_u(x_0))$$

where g was defined in (4.2).

Proof. We consider the same points x_0 as in Lemma 4.8. Take any infinitesimal sequence $(\varepsilon_j)_j$ such that

$$g(x_0, u^+(x_0), u^-(x_0), \nu_u(x_0)) = \lim_{j\to\infty} \frac{m(u_{x_0}, B_{\varepsilon_j}(x_0))}{2\varepsilon_j},$$

and recall that (4.28) is valid for $\mathcal{L}^1$-a.e. $s \in (0,1)$ (as usual $w_j = w_{\varepsilon_j}$). Having fixed such an s, let $z_j \in SBD^p(B_{s\,\varepsilon_j}(x_0))$ with $z_j = w_j$ on $\partial B_{s\varepsilon_j}(x_0)$ be such that

$$F(z_j, B_{s\varepsilon_j}(x_0)) \le m(w_j, B_{s\varepsilon_j}(x_0)) + \varepsilon_j^2.$$

Let $\varphi \in C_c^\infty(B_{\varepsilon_j}(x_0),[0,1])$ be a cut-off function such that $\varphi \equiv 1$ on $B_{s\varepsilon_j}(x_0)$ and $\|\nabla\varphi\|_{L^\infty} \le \frac{2}{(1-s)\varepsilon_j}$. Define $\zeta_j := \varphi\, z_j + (1-\varphi)u_{x_0}$, with the convention that z_j is extended equal to w_j outside $B_{s\varepsilon_j}(x_0)$. By using ζ_j as a test field for $m(u_{x_0}, B_{\varepsilon_j}(x_0))$ we infer from the growth condition in (1.1) and the locality of F

$$
\begin{aligned}
&m(u_{x_0}, B_{\varepsilon_j}(x_0)) \le F(\zeta_j, B_{\varepsilon_j}(x_0)) \le m(w_j, B_{s\varepsilon_j}(x_0)) + \varepsilon_j^2 \\
&+C\int_{B_{\varepsilon_j}(x_0)\setminus B_{s\,\varepsilon_j}(x_0)} (1+|e(w_j)|^p)\,dx + \frac{C}{((1-s)\varepsilon_j)^p}\int_{B_{\varepsilon_j}(x_0)\setminus B_{s\varepsilon_j}(x_0)} |w_j - u_{x_0}|^p dx \\
&\qquad\qquad + C\int_{(B_{\varepsilon_j}(x_0)\setminus B_{s\varepsilon_j}(x_0))\cap J_{\zeta_j}} (1+|[\zeta_j]|)d\mathcal{H}^1,
\end{aligned}
$$

where $C = C(\beta,p) > 0$. Arguing as in the corresponding estimate in Lemma 4.9 (cf. (4.29)), and by taking into account the choice of $(\varepsilon_j)_j$ we conclude that

$$
\begin{aligned}
g(x_0, u^+(x_0), u^-(x_0), \nu_u(x_0)) &\le \liminf_{j\to\infty} \frac{m(w_j, B_{s\varepsilon_j}(x_0))}{2\varepsilon_j} + C(1-s)(1+|[u](x_0)|) \\
&= s\frac{dF(u,\cdot)}{d\mathcal{H}^1 \llcorner J_u}(x_0) + C(1-s)(1+|[u](x_0)|).
\end{aligned}
$$

The last equality follows from (4.12). The conclusion is achieved by letting $s \uparrow 1$ in the last inequality, with $s \in (0,1)$ satisfying (4.28). □

4.4 Proof of Theorem 1.1

Proof of Theorem 1.1. The conclusion straightforwardly follows by Lemmata 4.6, 4.7, 4.9, and 4.10. □

Proposition 4.11. *The assertion in Theorem 1.1 holds also if property (iv) is replaced by the weaker*

(iv') There are $\alpha, \beta > 0$ such that for any $u \in SBD^p(\Omega)$, any $B \in \mathcal{B}(\Omega)$,

$$
\begin{aligned}
&\alpha\Big(\int_B |e(u)|^p dx + \mathcal{H}^1(J_u \cap B)\Big) \le F(u,B) \\
&\le \beta\Big(\int_B (|e(u)|^p + 1)dx + \int_{J_u\cap B} (1+|[u]|)d\mathcal{H}^1\Big).
\end{aligned}
$$

Proof. Given F satisfying properties (i)-(iii) and (iv'), we define for $\delta > 0$ a functional $F_\delta : SBD^p(\Omega)\times\mathcal{B}(\Omega) \to [0,\infty)$ by

$$
F_\delta(u,B) := F(u,B) + \delta\int_{J_u\cap B} |[u]|d\mathcal{H}^1,
$$

for $u \in SBD^p(\Omega)$ and $B \in \mathcal{B}(\Omega)$. Since F_δ satisfies properties (i)-(iv), there are two functions f and g_δ such that F_δ can be represented as in (1.2). The family of functionals F_δ is pointwise increasing in δ, therefore there exists the pointwise limit g of g_δ as $\delta \to 0$. We conclude that the representation (1.2) holds for F with densities f and g. □

Remark 4.12. *Since F is lower semicontinuous on $W^{1,p}$, the integrand f is quasiconvex [1, 36]. Since F is lower semicontinuous on piecewise constant functions, g is BV-elliptic [2, 3].*

Remark 4.13. *If the functional F additionally obeys*

$$F(u + I, B) = F(I, B),$$

for every $u \in SBD^p(\Omega)$, every ball $B \subset \Omega$, and every affine function I such that $e(I) = 0$, then there are two functions $f : \Omega \times \mathbb{R}^{2\times 2} \to [0, \infty)$ and $g : \Omega \times \mathbb{R}^2 \times S^1 \to [0, \infty)$ such that

$$F(u, B) = \int_B f(x, e(u(x)))dx + \int_{B\cap J_u} g(x, [u](x), \nu_u(x))d\mathcal{H}^1 .$$

Remark 4.14. *A growth condition on the volume part of the type of (1.1) alone does not force the energy density to depend only on $e(u)$. As an example, the integrand $f : \mathbb{R}^{2\times 2} \to [0, \infty)$ defined by*

$$f(\xi) := (\xi_{11} + \xi_{22})^2 + \sqrt{(\xi_{12}^2 + \xi_{21}^2)^2 + 1} - 2\det(\xi)$$

satisfies

$$\frac{1}{8}|\xi + \xi^T|^2 \le f(\xi) \le \frac{1}{4}|\xi + \xi^T|^2 + 1$$

for every $\xi \in \mathbb{R}^{2\times 2}$, but evidently $f(\xi)$ depends also on the skew-symmetric part $\xi - \xi^T$. At the same time, f is quasiconvex. We do not know if there is g such that the functional F defined as in (1.2) satisfies the growth condition (1.1) and is lower semicontinuous.

Acknowledgments

F. Iurlano wishes to thank Gianni Dal Maso for an interesting discussion. This work was partially supported by the Deutsche Forschungsgemeinschaft through the Sonderforschungsbereich 1060 *"The mathematics of emergent effects"*, project A6. S. Conti thanks the University of Florence for the warm hospitality of the DiMaI "Ulisse Dini", where part of this work was carried out. M. Focardi and F. Iurlano are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

References

[1] E. Acerbi and N. Fusco, *Semicontinuity problems in the calculus of variations*, Arch. Rat. Mech. Anal., 86 (1984), pp. 125–145.

[2] L. Ambrosio and A. Braides, *Functionals defined on partitions in sets of finite perimeter. I. Integral representation and* Γ*-convergence*, J. Math. Pures Appl. (9), 69 (1990), pp. 285–305.

[3] ——, *Functionals defined on partitions in sets of finite perimeter. II. Semicontinuity, relaxation and homogenization*, J. Math. Pures Appl. (9), 69 (1990), pp. 307–333.

[4] L. Ambrosio, A. Coscia, and G. Dal Maso, *Fine properties of functions with bounded deformation*, Arch. Rational Mech. Anal., 139 (1997), pp. 201–238.

[5] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of bounded variation and free discontinuity problems*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 2000.

[6] G. Anzellotti and M. Giaquinta, *Existence of the displacement field for an elastoplastic body subject to Hencky's law and von Mises yield condition*, Manuscripta Math., 32 (1980), pp. 101–136.

[7] G. Bellettini, A. Coscia, and G. Dal Maso, *Compactness and lower semicontinuity properties in* $SBD(\Omega)$, Math. Z., 228 (1998), pp. 337–351.

[8] G. Bouchitté and G. Dal Maso, *Integral representation and relaxation of convex local functionals on* $\mathrm{BV}(\Omega)$, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 20 (1993), pp. 483–533.

[9] G. Bouchitté, I. Fonseca, G. Leoni, and L. Mascarenhas, *A global method for relaxation in* $W^{1,p}$ *and in* SBV_p, Arch. Ration. Mech. Anal., 165 (2002), pp. 187–242.

[10] G. Bouchitté, I. Fonseca, and L. Mascarenhas, *A global method for relaxation*, Arch. Rational Mech. Anal., 145 (1998), pp. 51–98.

[11] A. Braides and V. Chiadò Piat, *Integral representation results for functionals defined on* $\mathrm{SBV}(\Omega;\mathbf{R}^m)$, J. Math. Pures Appl. (9), 75 (1996), pp. 595–626.

[12] G. Buttazzo and G. Dal Maso, Γ-*limits of integral functionals*, J. Analyse Math., 37 (1980), pp. 145–185.

[13] ——, *A characterization of nonlinear functionals on Sobolev spaces which admit an integral representation with a Carathéodory integrand*, J. Math. Pures Appl. (9), 64 (1985), pp. 337–361.

[14] ——, *Integral representation and relaxation of local functionals*, Nonlinear Anal., 9 (1985), pp. 515–532.

[15] L. Carbone and C. Sbordone, *Some properties of* Γ*-limits of integral functionals*, Ann. Mat. Pura Appl. (4), 122 (1979), pp. 1–60.

[16] A. Chambolle, *An approximation result for special functions with bounded deformation*, J. Math. Pures Appl. (9), 83 (2004), pp. 929–954.

[17] ——, *Addendum to: "An approximation result for special functions with bounded deformation" [16]*, J. Math. Pures Appl. (9), 84 (2005), pp. 137–145.

[18] A. Chambolle, S. Conti, and G. Francfort, *Korn-Poincaré inequalities for functions with a small jump set*, Preprint hal-01091710v1, (2014).

[19] A. Chambolle, A. Giacomini, and M. Ponsiglione, *Piecewise rigidity*, J. Funct. Anal., 244 (2007), pp. 134–153.

[20] S. Conti, M. Focardi, and F. Iurlano, *Existence theorem for the Griffith fracture model in dimension two*, In preparation.

[21] S. Conti, M. Focardi, and F. Iurlano, *Which special functions of bounded deformation have bounded variation?*, Preprint arXiv:1502.07464, (2015).

[22] S. Conti and B. Schweizer, *Rigidity and gamma convergence for solid-solid phase transitions with SO(2) invariance*, Comm. Pure Appl. Math., 59 (2006), pp. 830–868.

[23] ——, *A sharp-interface limit for a two-well problem in geometrically linear elasticity*, Arch. Ration. Mech. Anal., 179 (2006), pp. 413–452.

[24] G. Dal Maso, *Integral representation on* BV(Ω) *of* Γ*-limits of variational integrals*, Manuscripta Math., 30 (1979/80), pp. 387–416.

[25] ——, *An introduction to* Γ*-convergence*, Progress in Nonlinear Differential Equations and their Applications, 8, Birkhäuser Boston Inc., Boston, MA, 1993.

[26] ——, *Generalised functions of bounded deformation*, J. Eur. Math. Soc. (JEMS), 15 (2013), pp. 1943–1997.

[27] G. Dal Maso and L. Modica, *A general theory of variational functionals*, in Topics in functional analysis, 1980–81, Quaderni, Scuola Norm. Sup. Pisa, Pisa, 1981, pp. 149–221.

[28] ——, *Integral functionals determined by their minima*, Rend. Sem. Mat. Univ. Padova, 76 (1986), pp. 255–267.

[29] E. De Giorgi, *Sulla convergenza di alcune successioni d'integrali del tipo dell'area*, Rend. Mat. (6), 8 (1975), pp. 277–294. Collection of articles dedicated to Mauro Picone on the occasion of his ninetieth birthday.

[30] E. De Giorgi, M. Carriero, and A. Leaci, *Existence theorem for a minimum problem with free discontinuity set*, Arch. Rational Mech. Anal., 108 (1989), pp. 195–218.

[31] F. Ebobisse and R. Toader, *A note on the integral representation representation of functionals in the space* $SBD(\Omega)$, Rend. Mat. Appl. (7), 23 (2003), pp. 189–201 (2004).

[32] M. Friedrich, *A Korn-Poincaré-type inequality for special functions of bounded deformation*, preprint, (2015).

[33] ——, *A Korn-type inequality in* SBD *for functions with small jump sets*, preprint, (2015).

[34] G. Friesecke, R. D. James, and S. Müller, *A theorem on geometric rigidity and the derivation of nonlinear plate theory from three-dimensional elasticity*, Comm. Pure Appl. Math., 55 (2002), pp. 1461–1506.

[35] R. Kohn and R. Temam, *Dual spaces of stresses and strains, with applications to Hencky plasticity*, Appl. Math. Optim., 10 (1983), pp. 1–35.

[36] P. Marcellini, *Approximation of quasiconvex functions, and lower semicontinuity of multiple integrals*, Manuscripta Math., 51 (1985), pp. 1–28.

[37] J. A. Nitsche, *On Korn's second inequality*, RAIRO Anal. Numér., 15 (1981), pp. 237–248.

[38] C. Sbordone, *Su alcune applicazioni di un tipo di convergenza variazionale*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 2 (1975), pp. 617–638.

[39] P.-M. Suquet, *Sur un nouveau cadre fonctionnel pour les équations de la plasticité*, C. R. Acad. Sci. Paris Sér. A-B, 286 (1978), pp. A1129–A1132.

[40] R. Temam, *Problèmes mathématiques en plasticité*, vol. 12 of Méthodes Mathématiques de l'Informatique, Gauthier-Villars, Montrouge, 1983.

[41] R. Temam and G. Strang, *Functions of bounded deformation*, Arch. Rational Mech. Anal., 75 (1980/81), pp. 7–21.